

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number
WO 02/24698 A1

(51) International Patent Classification⁷: **C07D 473/04**,
473/06, 473/08, A61K 31/522, A61P 15/00, 9/00

(21) International Application Number: PCT/US01/28983

(22) International Filing Date:
17 September 2001 (17.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/233,567 19 September 2000 (19.09.2000) US

(71) Applicant: **SCHERING CORPORATION** [US/US];
Patent Department - K-6-1 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: **CHACKALAMANNIL, Samuel**; 79 Strat-
ford Road, East Brunswick, NJ 08816 (US). **WANG,**
Yuguang; 38 Princess Drive, North Brunswick, NJ 08902
(US). **BOYLE, Craig, D.**; 13 Mill Lane, Branchburg, NJ
08876 (US). **STAMFORD, Andrew, W.**; 27 Overlook
Road, Chatham Township, NJ 07928 (US).

(74) Agent: **KUTZENCO, Allan, N.**; Schering Corporation,
Patent Department - K-6-1 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT,
RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ,
VN, YU, ZA.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: XANTHINE PHOSPHODIESTERASE V INHIBITORS

(57) Abstract: A xanthine phosphodiesterase V inhibitor having the formula (I), where R⁴ is a C₃₋₁₅ cycloalkyl group, with or without one or more substituents, C₃₋₁₅ cycloalkenyl group, with or without one or more substituents, or a heterocycloalkyl group of 3 to 15 members, with or without one or more substituents; useful for treating male (erectile) and female sexual dysfunction and other physiological disorders. For example, a representative compound of the invention is formula (II).



WO 02/24698 A1

TITLE OF THE INVENTION

Xanthine Phosphodiesterase V Inhibitors

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

10

The invention relates to polycyclic nucleotide xanthine phosphodiesterase V inhibitors.

2. Description of Related Art

15

Phosphodiesterase ("PDE") V inhibitor compounds are described by Kenneth J. Murray in *Phosphodiesterase V_A Inhibitors*, DN & P 6(3), pp. 150-156 (April, 1993), which is hereby incorporated herein by reference in its entirety, to have potential therapeutic value for a number of physiological disorders. One compound disclosed in the Murray article is MIMAX, a polycyclic xanthine PDE V inhibitor substituted at its 8-position with a -NHCH₃ group.

20

US 5,409,934, which is hereby incorporated herein by reference in its entirety, discloses a series of xanthine PDE V inhibitors that are substituted at the 8-position with, among other possibilities, one of the following groups: -NO₂, -NR^sR^t or -NR⁶SO₂R⁵, where R^s and R^t, independently of one another, are each a hydrogen atom or an alkyl group, or R^s and R^t, together with the nitrogen atom to which they are both attached, form a phthalimido group, R⁵ is

25

an alkyl or aryl group, and R^6 is a hydrogen atom or $-SO_2R^7$, where R^7 is an alkyl or aryl group.

US 5,470,579, which is hereby incorporated herein by reference in its entirety, discloses a xanthine PDE V inhibitor having a substituted or
5 unsubstituted $-NH_2$ group at the 8-position, for example, $-NHR$, where R is a C_1 - C_6 alkyl group.

WO 93/23401, which is hereby incorporated herein by reference in its entirety, discloses xanthine PDE V inhibitors that are substituted at the 8-position with $-NH(CH_2)_2CH(CH_2OR^4)_2$.

10 WO 92/05176, which is hereby incorporated herein by reference in its entirety, discloses 8-acylaminoxanthine PDE V inhibitors that are substituted at the 8-position with $-NHCOC_6H_5COOH$.

WO 92/05175, which is hereby incorporated herein by reference in its entirety, discloses 8-aminoxanthine PDE V inhibitors that are substituted at
15 the 8-position with $-NH_2$ or $-NHR$, where R is an alkyl, arylalkyl or unsaturated heterocyclic (e.g., heteroaryl) group.

Specific PDE V inhibitors have been found useful for specific indications. For example, the use of PDE V inhibitors for treating impotence has met with commercial success with the introduction of sildenafil citrate,
20 better known as Viagra® (Pfizer, NY, NY). The chemistry and use of Viagra®, including its mechanism of action in treating erectile dysfunction, are taught in EP 0 702 555 B1, which is hereby incorporated herein by reference in its entirety. Additional PDE V inhibitors useful for treating erectile dysfunction

are disclosed in WO 99/24433, which is hereby incorporated herein by reference in its entirety.

Erectile dysfunction is a treatable and highly recognized health concern, affecting more than 30 million men in the United States, including one in four over age 65. Erectile dysfunction occurs when a man consistently is unable to sustain an erection sufficient for conducting sexual intercourse. In the past, psychological reasons were the most common explanation for erectile dysfunction or it was considered a natural part of aging. However, researchers today acknowledge that more than 70 percent of instances of erectile dysfunction are due to physical or medical problems. There are several factors that may contribute to erectile dysfunction, including:

- Poor blood circulation – atherosclerosis or hardening of the arteries, high blood pressure and high cholesterol.
- Neurological disorders – multiple sclerosis, Alzheimer's disease and Parkinson's disease.
- Hormone imbalances – diabetes, thyroid disorders and low testosterone levels.
- Trauma – spinal cord injury, prostate surgery or other trauma to the pelvic area.
- Prescription and over-the-counter medications – blood pressure medications, antidepressants and certain drug combinations.
- Lifestyle habits – smoking, alcohol abuse and using illegal drugs.

U.S. 5,939,419 and U.S. 5,393,755, both of which are hereby incorporated herein by reference in their entirety, disclose polycyclic guanine PDE V derivatives that are useful for the treatment of cardiovascular and pulmonary disorders.

5 As has been shown by the representative art cited above, certain xanthine/guanine PDE V inhibitors have been found to be useful for treating cardiovascular and pulmonary disorders, while some others have been found useful for treating impotence. It has been further shown that certain xanthine PDE V inhibitors can be substituted at the 8-position by a variety of groups,
10 including nitro and unsubstituted or substituted amino groups. The substituted amino groups include saturated heterocycles, where the nitrogen atom and its substituents together form an unsaturated heterocyclic group (e.g., -NR^xR^y can form a heterocycle).

 It is an object of this invention to provide a polycyclic xanthine PDE V
15 inhibitor that possesses beneficial therapeutic properties.

 It is a further object of the invention to provide a polycyclic xanthine PDE V inhibitor that has especially useful pharmacological properties.

 It is yet another object of the invention to provide a polycyclic xanthine PDE V inhibitor that has good metabolic stability.

20 It is still another object of the invention to provide a polycyclic xanthine PDE V inhibitor that is effective for treating a variety of physiological symptoms and diseases in which PDE V plays a role.

It is also an object of the invention to provide a polycyclic xanthine PDE V inhibitor that is especially effective for treating erectile dysfunction with minimal side effects.

These and other objects of the invention will become apparent as the description progresses.

Definitions and Usage of Terms

The following definitions and terms are used herein or are otherwise known to a skilled artisan. Except where stated otherwise, the following definitions apply throughout the specification and claims. These definitions apply regardless of whether a term is used by itself or in combination with other terms, unless otherwise indicated. Hence, the definition of "alkyl" applies to "alkyl" as well as the "alkyl" portions of "hydroxyalkyl," "haloalkyl," "alkoxy," etc.


The term "chemically-compatible," as used herein, means that a substituent or variable in a structure, process or the like is selected to be capable of resulting in a stable compound.

The term "substituted" or the phrase "with . . . one or more substituents," as used herein, means the replacement of one or more atoms or radicals, usually hydrogen atoms, in a given structure with a chemically-compatible atom(s) or radical(s) selected from a specified group. In the situations where more than one atom or radical may be replaced with substituents selected from the same specified group, the substituents may be, unless otherwise specified, either the same or different at every position.

Radicals of specified groups, such as alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, arylalkyl, alkylaryl, heterocycloalkyl, aryl and heteroaryl groups, independently of or together with one another, may be substituents for any substituted group, unless otherwise known, stated or shown to be to the contrary.

Representative substituents for alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, arylalkyl, alkylaryl, aryl, heteroaryl and heterocycloalkyl groups include, but are not limited to, the following moieties: alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, arylalkyl, alkylaryl, aryl, heteroaryl, heterocycloalkyl, hydroxyalkyl, arylalkyl, aminoalkyl, haloalkyl, thioalkyl, alkylthioalkyl, carboxyalkyl, imidazolylalkyl, indolylalkyl, mono-, di- and trihaloalkyl, mono-, di- and trihaloalkoxy, amino, alkylamino, dialkylamino, alkoxy, hydroxy, halo (e.g., -Cl and -Br), nitro, oximino, -COOR⁵⁰, -COR⁵⁰, -SO₀₋₂R⁵⁰, -SO₂NR⁵⁰R⁵¹, NR⁵²SO₂R⁵⁰, =C(R⁵⁰R⁵¹), =N-OR⁵⁰, =N-CN, =C(halo)₂, =S, =O, -CON(R⁵⁰R⁵¹), -OCOR⁵⁰, -OCON(R⁵⁰R⁵¹), -N(R⁵²)CO(R⁵⁰), -N(R⁵²)COOR⁵⁰ and -N(R⁵²)CON(R⁵⁰R⁵¹), where:

R⁵⁰, R⁵¹ and R⁵² may be independently selected from the following: a hydrogen atom and a branched or straight-chain, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₄₋₆ heterocycloalkyl, heteroaryl and aryl group, with or without substituents. When permissible, R⁵⁰ and R⁵¹ can be joined together to form a carbocyclic or heterocyclic ring system. R⁵⁰, R⁵¹ and R⁵² may also include:



where,

5 R^{40} and R^{41} are, independently of one another, each a hydrogen atom or a branched or straight-chain, optionally substituted, alkyl, cycloalkyl, heterocycloalkyl, halo, aryl, imidazolylalkyl, indolylalkyl,

heteroaryl, arylalkyl, arylalkoxy, heteroarylalkyl, heteroarylalkoxy, aminoalkyl, haloalkyl, mono-, di- or trihaloalkyl, mono-, di- or trihaloalkoxy, nitro, cyano, alkoxy, hydroxy, amino, phosphino, phosphate, alkylamino, dialkylamino, formyl, alkylthio, trialkylsilyl, alkylsulfonyl, arylsulfonyl, alkylsulfinyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, hydroxyalkyl, morpholino, thioalkyl, alkylthioalkyl, carboxyalkyl, oximino, $-\text{COOR}^{50}$, $-\text{COR}^{50}$, $-\text{SO}_{0-2}\text{R}^{50}$, $-\text{SO}_2\text{NR}^{50}\text{R}^{51}$, $-\text{NR}^{52}\text{SO}_2\text{R}^{50}$, $-\text{CON}(\text{R}^{50}\text{R}^{51})$, $-\text{OCON}(\text{R}^{50}\text{R}^{51})$, $-\text{N}(\text{R}^{52})\text{CO}(\text{R}^{50})$, $-\text{N}(\text{R}^{52})\text{COOR}^{50}$, $-\text{N}(\text{R}^{52})\text{CON}(\text{R}^{50}\text{R}^{51})$ or $-\text{OCONR}^{50}$ group, where, R^{50} , R^{51} and R^{52} are as defined above;

R^{42} is a hydrogen atom or a branched or straight-chain, optionally substituted, alkyl, alkenyl, arylalkyl or acyl group; and

R^{43} is a hydrogen atom or a branched or straight-chain, optionally substituted, alkyl or aryl group;

wherein, the optional substituents are defined the same as above for the one or more substituents.

Preferred substituents on aryl and heteroaryl groups include, but are not limited to, any of the moieties recited above in the definition for R^{40} and R^{41} .

The term "heteroatom," as used herein, means a nitrogen, sulfur, or oxygen atom. Multiple heteroatoms in the same group may be the same or different.

The term "hydrocarbon," as used herein, means a compound or radical consisting of only carbon and hydrogen atoms, including aliphatic, aromatic, normal, saturated and unsaturated hydrocarbons.

5 The term "alkyl," as used herein, means an unsubstituted or substituted, straight or branched, hydrocarbon chain (*i.e.*, comprising carbon and hydrogen atoms bonded together), having, preferably, from one to twenty-four carbon atoms, more preferably, from one to twelve carbon atoms, and most preferably, from one to eight carbon atoms.

10 The term "cycloalkyl" or "cycloalkane," as used herein, means an unsubstituted or substituted, saturated, stable non-aromatic carbocyclic ring, having, preferably, from three to fifteen carbon atoms, more preferably, from three to eight carbon atoms. The carbon ring radical is saturated and may be fused, for example, benzofused, with one to three cycloalkyl, aromatic, heterocyclic or heteroaromatic rings. The cycloalkyl may be attached at any
15 endocyclic carbon atom that results in a stable structure. Preferred carbocycles have from five to six carbons. Examples of carbocycle radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like.

20 The term "alkenyl," as used herein, means an unsubstituted or substituted, unsaturated, straight or branched, hydrocarbon chain having at least one double bond present and, preferably, from two to fifteen carbon atoms, more preferably, from two to twelve carbon atoms.

The term "cycloalkenyl," as used herein, means an unsubstituted or substituted, unsaturated carbocyclic ring having at least one double bond

present and, preferably, from three to fifteen carbon atoms, more preferably, from five to eight carbon atoms. A cycloalkenyl group is an unsaturated carbocyclic group. Examples of cycloalkenyl groups include cyclopentenyl and cyclohexenyl.

5 The term "alkynyl," as used herein, means an unsubstituted or substituted, unsaturated, straight or branched, hydrocarbon chain having at least one triple bond present and, preferably, from two to twelve carbon atoms, more preferably, two to ten carbon atoms.

10 The term "bicycloalkyl," as used herein, represents a saturated linearly fused or bridged carbocyclic ring having, preferably, from 5 to 12 carbon atoms.

15 The term "aryl," as used herein, means a substituted or unsubstituted, aromatic, mono- or bicyclic carbocyclic ring system having from one to two aromatic rings. The aryl moiety will generally have from 6 to 14 carbon atoms with all available substitutable carbon atoms of the aryl moiety being intended as possible points of attachment. Representative examples include phenyl, tolyl, xylyl, cumenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like. If desired, the carbocyclic moiety can be substituted with from one to five, preferably, one to three moieties, such as mono- through pentahalo, 20 alkyl, trifluoromethyl, phenyl, hydroxy, alkoxy, phenoxy, amino, monoalkylamino, dialkylamino and the like.

 The term "heteroaryl," as used herein, means a mono- or bicyclic ring system containing one or two aromatic rings and at least one nitrogen, oxygen or sulfur atom in an aromatic ring. Heteroaryl groups (including bicyclic

heteroaryl groups) can be unsubstituted or substituted with a plurality of substituents, preferably, one to five substituents, more preferably, one, two or three substituents (e.g., mono- through pentahalo, alkyl, trifluoromethyl, phenyl, hydroxy, alkoxy, phenoxy, amino, monoalkylamino, dialkylamino and the like). Typically, a heteroaryl group represents a cyclic group of five or six atoms, or a bicyclic group of nine or ten atoms, at least one of which is carbon, and having at least one oxygen, sulfur or nitrogen atom interrupting a carbocyclic ring having a sufficient number of π (π) electrons to provide aromatic character. Representative heteroaryl (heteroaromatic) groups are pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, furanyl, benzofuranyl, thienyl, benzothienyl, thiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, triazolyl, isothiazolyl, benzothiazolyl, benzoxazolyl, oxazolyl, pyrrolyl, isoxazolyl, 1,3,5-triazinyl and indolyl groups.

The term "arylalkyl," as used herein, means an alkyl moiety substituted with an optionally substituted, aryl or heteroaryl group. Representative arylalkyl groups include a benzyl group and fused bicyclic systems which contain one aryl group.

The term "alkylaryl," as used herein, means an aryl or heteroaryl moiety substituted with an optionally substituted, alkyl group. Representative alkylaryl groups include o-, m- and p-linked tolyl and xylyl groups.

Unless otherwise known, stated or shown to be to the contrary, the point of attachment for a multiple term substituent (multiple terms that are combined to identify a single moiety) to a subject structure is through the last named term of the multiple term. For example, an "arylalkyl" substituent

attaches to a targeted structure through the "alkyl" portion of the substituent. Conversely, when the substituent is "alkylaryl", it attaches to a targeted structure through the "aryl" portion of the substituent. Similarly, a cycloalkylalkyl substituent attaches to a targeted through the latter "alkyl" portion of the substituent (e.g., Structure-alkyl-cycloalkyl).

The term "heterocycloalkyl," as used herein, means an unsubstituted or substituted, saturated cyclic ring system having from three to fifteen members, preferably, from three to eight members, and comprising carbon atoms and at least one heteroatom as part of the ring.

The term "heterocyclic ring" or "heterocycle," as used herein, means an unsubstituted or substituted, saturated, unsaturated or aromatic ring, comprised of carbon atoms and one or more heteroatoms in the ring. Heterocyclic rings may be monocyclic or polycyclic. Monocyclic rings preferably contain from three to eight atoms, most preferably, five to seven atoms. Polycyclic ring systems consisting of two rings preferably contain from six to sixteen atoms, most preferably, ten to twelve atoms. Polycyclic ring systems consisting of three rings contain, preferably, from thirteen to seventeen atoms, most preferably, fourteen to fifteen atoms. Each heterocyclic ring has at least one hetero atom. Unless otherwise stated, the heteroatoms may be independently selected from the following: nitrogen, sulfur and oxygen atoms.

The term "carbocyclic ring" or "carbocycle," as used herein, means an unsubstituted or substituted, saturated, unsaturated or aromatic (e.g., aryl), hydrocarbon ring, unless otherwise specifically identified. Carbocycles may

be monocyclic or polycyclic. Monocyclic rings preferably contain from three to eight atoms, most preferably, five to seven atoms. Polycyclic rings having two rings preferably contain from six to sixteen atoms, most preferably, ten to twelve atoms, and those having three rings preferably contain from thirteen to
5 seventeen atoms, most preferably, fourteen to fifteen atoms.

The term "alkoxy," as used herein, means an oxygen atom bonded to a hydrocarbon chain, such as an alkyl or alkenyl group (e.g., -O-alkyl or -O-alkenyl). Representative alkoxy groups include methoxy, ethoxy, and isopropoxy groups.

10 The term "hydroxyalkyl," as used herein, means a substituted hydrocarbon chain, preferably, an alkyl group, having at least one hydroxy substituent (i.e., -OH). Additional substituents to the alkyl group may also be present. Representative hydroxyalkyl groups include hydroxymethyl, hydroxyethyl and hydroxypropyl groups.

15 The term "carboxyalkyl," as used herein, means a substituted hydrocarbon chain, preferably, a substituted alkyl group, which has a carboxyl substituent (e.g., -COOH) and may also have additional substituents (such as one of the representative substituents identified above for the term "substituted"). Representative carboxyalkyl groups include carboxymethyl (-
20 CH₂CO₂H) and carboxyethyl (-CH₂CH₂CO₂H) groups, and derivatives thereof, such as the corresponding esters.

The term "aminoalkyl," as used herein, means an alkyl group substituted with an amine moiety (e.g., -alkylNH₂), such as aminomethyl.

The term "alkylamino," as used herein, means an amino moiety having from one or two alkyl substituents (e.g., -NH-alkyl), such as dimethylamino.

5 The term "alkenylamino," as used herein, means an amino moiety having from one or two alkenyl substituents, where the nitrogen atom of the amino group is not attached to the alkene-forming carbon atom (e.g., -NH-CH₂-alkenyl), such as dibutenylamino.

The term "arylamino," as used herein, means an amine moiety substituted with an aryl group (i.e., -NH-aryl).

10 The term "alkylimino," as used herein, means an imino moiety having one alkenyl or two alkyl substituents (e.g., -C=N-alkyl).

The term "oximino," as used herein, means compounds containing the -C=N-OR⁶⁹ radical, where R⁶⁹ is a hydrogen atom or an alkyl or aryl group.

The term "aroyl," as used herein, means the radical R-CO-; where R is an aromatic group. Representative aroyls are benzoyl and naphthoyl.

15 The term "aryloxy," as used herein, means an oxygen atom having an aryl substituent (e.g., -O-aryl).

The term "ester," as used herein, means compounds containing a substituted carboxylic acid (e.g., -COO-aryl).

20 The term "acyl" or "carbonyl," as used herein, means a carbon to oxygen double bond, (e.g., R-C(=O)-), which can be a radical of a carboxylic acid having the formula alkyl-CO-, aryl-CO-, arylalkyl-CO-, cycloalkyl-CO-, alkylcycloalkyl-CO- or heteroaryl-CO-. Representative acyl groups include acetyl, propionyl, butanoyl and benzoyl groups.

The term "acyloxy," as used herein, means an oxygen atom having an acyl substituent (*e.g.*, -O-acyl), for example, -O-C(=O)-alkyl.

5 The term "acylamino," as used herein, means an amino moiety having an acyl substituent (*e.g.*, -NH-acyl), for example, an amide with the formula -NH-(C=O)-alkyl, a urea with the formula -NH-(C=O)-NH-alkyl or a carbamate with the formula -NH-(C=O)-OR, where R is an alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, arylalkyl or heterocycloalkyl group.

10 The term "halo," "halogen" or "halide," as used herein, means a chloro, bromo, fluoro or iodo atom radical. Chlorides, bromides and fluorides are preferred halides.

The term "lower hydrocarbon" (*e.g.*, "lower alkyl"), as used herein, means a hydrocarbon chain comprised of from, unless otherwise stated, one to eight carbon atoms, preferably, one to six carbon atoms, and most preferably, one to four carbon atoms.

15 The term "polyhalo," as used herein, represents substitution of at least two halo atoms to a group modified by the term "polyhalo."

The term "aminosulfonyl," as used herein, represents a group having the formula: -SO₂NR⁷⁹R⁸⁹, where R⁷⁹ and R⁸⁹ are, independently of one another, each a hydrogen atom or a lower alkyl (*e.g.*, from 1 to 6 carbon atoms) or aryl group.

20

The term "sulfonyl," as used herein, represents a group having the formula: -S(O)₂-.

When a variable appears more than once in a structural formula, for example, R⁵⁹ for where X is -C(OR⁵⁹)₂-, the identity of each variable appearing

more than once may be independently selected from the definition for that variable.

The term "prodrug," as used herein, represents a compound that is a drug precursor, which following administration to a patient, releases a drug *in vivo* via some kind of chemical and/or physiological process (e.g., a prodrug on being brought to a physiological *pH* and/or through an enzyme action is converted to a desired drug form).

The term "compound of the formula (I.1) or (II.1)", as used herein, represents a compound having a chemical structure encompassed by the formula (I.1) or (II.1), and includes any and all enantiomers, stereoisomers, rotomers, tautomers and prodrugs of the compound. Compounds of the formula (I.1) or (II.1) also include their corresponding pharmaceutically-acceptable salts, solvates, esters and derivatives.

The term "pharmaceutically-acceptable excipients," as used herein, includes any physiologically inert, pharmacologically inactive material known to one skilled in the art, which is compatible with the physical and chemical characteristics of the particular active ingredient selected for use.

Pharmaceutically-acceptable excipients include polymers, resins, plasticizers, fillers, binders, lubricants, glidants, disintegrates, solvents, co-solvents, buffer systems, surfactants, preservatives, sweetening agents, flavoring agents, pharmaceutical grade dyes or pigments, and viscosity agents.

The term "pharmaceutical composition," as used herein, means a combination of at least one inventive compound (e.g., PDE V inhibitor) and at least one pharmaceutically-acceptable excipient.

The terms "compound [having the formula (I)] or a pharmaceutical composition thereof" include neutral, acidic and alkaline forms of the compound or composition, as well as solvates, esters and salts (as are defined below) thereof, and further includes derivatives of the inventive compounds.

The term "pharmaceutically-acceptable salt," as used herein, means a cationic salt formed at an acidic (e.g., carboxyl) group or an anionic salt formed at a basic (e.g., amino) group of the compound. Many such salts are known in the art, for example, those that are described in WO 87/05297 (1987), which is hereby incorporated in its entirety by reference herein. Preferred cationic salts include the alkali-metal salts (e.g., sodium and potassium) and alkaline earth metal salts (e.g., magnesium and calcium). Preferred anionic salts include the halide (e.g., chloride), acetate and phosphate salts.

The phrase "effective amount," as used herein, means an amount of a compound or composition which is sufficient to significantly and positively modify the symptoms and/or conditions to be treated (e.g., provide a positive clinical response). The phrase "safe and effective amount," as used herein, means that an "effective amount" must also be safe, that is, an amount that is sufficient to provoke a positive response, yet is small enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. The effective amount of an active ingredient for use in a pharmaceutical composition will vary with the particular condition being treated, the severity of the condition, the duration of the treatment, the nature

of concurrent therapy, the particular active ingredient being employed, the particular pharmaceutically-acceptable excipients utilized and like factors within the knowledge and expertise of the attending physician.

5 The phrase "administering [to a patient a safe and effective amount of the inventive compound]," as used herein, refers to any mode of introducing any form (e.g., solid, liquid or gas) of the inventive compounds *in vivo* to a patient (e.g., human or mammal). For example, introduction of the inventive compound to a patient may be accomplished via oral ingestion (e.g., tablets, capsules, gels, solutions, etc.), adsorption, absorption (e.g., transmucosal
10 sublingual or buccal administration), transdermal applications (e.g., topical applications via patches, lotions, etc.), suppositories, etc.

The term "oral dosage form," as used herein, means any pharmaceutical composition intended to be systemically administered to an individual by delivering the composition to the gastrointestinal tract of an
15 individual, via the mouth of the individual. For purposes of the invention, the delivered form can be a tablet (coated or non-coated), solution, suspension or capsule (coated or non-coated).


20 The term "injection," as used herein, means any pharmaceutical composition intended to be systemically administered to a human or other mammal, via delivery of a solution or emulsion containing the active ingredient, by puncturing the skin of said individual, in order to deliver the solution or emulsion to the circulatory system of the individual either by intravenous, intramuscular, intraperitoneal or subcutaneous injection.

Other than as shown in the operating examples or where is otherwise indicated, all numbers used in the specification and claims expressing quantities of ingredients, reaction conditions, and so forth, are understood as being modified in all instances by the term "about."

5

SUMMARY OF THE INVENTION

The invention comprises a compound having the formula (I):



(I)

where,

- (a) R¹ and R² are, independently of one another, each a C₁₋₁₅ alkyl group, branched or straight chain, with or without one or more substituents, such as a hydroxy or alkoxy substituent group, a C₂₋₁₅ alkenyl group, branched or straight chain, with or without one or more substituents, a C₂₋₁₅ alkynyl group, branched or straight chain, with or without one or more substituents, a C₃₋₁₅ cycloalkyl group, with or without one or more substituents, an arylalkyl group, with or without one or more substituents, an aryl group, with or without one or more

10



15

20

substituents, a heteroaryl group, with or without one or more substituents, $-OR^5$, $-COOR^5$, $-C(O)R^5$ or $-C(O)N(R^5)_2$, where R^5 is a hydrogen atom or a hydrocarbon radical, with or without one or more substituents, preferably, R^5 is a hydrogen atom or an alkyl group, branched or straight chain, with or without one or more substituents; or

one of R^1 and R^2 is equal to a hydrogen atom, and the other one of R^1 and R^2 is defined the same as above;

- 10 (b) R^3 is an aryl group, with or without one or more substituents, such as a hydroxy or alkoxy substituent group, a heteroaryl group, with or without one or more substituents, or a heterocyclic group having from 1 to 3 heteroatoms fused to a 5- or 6-membered aryl ring, with or without one or more
- 15 substituents, with the proviso that R^3 is not an aryl group substituted at its para position with a $-Y$ -aryl group, where Y is a carbon-carbon single bond, $-CO-$, $-O-$, $-S-$, $-N(R^{21})-$, $-CON(R^{22})-$, $-N(R^{22})CO-$, $-OCH_2-$, $-CH_2O-$, $-SCH_2-$, $-CH_2S-$, $-NHC(R^{23})(R^{24})-$, $-NR^{23}SO_2-$, $-SO_2NR^{23}-$, $-C(R^{23})(R^{24})NH-$, $-CH=CH-$, $-CF=CF-$,
- 20 $-CH=CF-$, $-CF=CH-$, $-CH_2CH_2-$, $-CF_2CF_2-$,



where,

R^{21} is a hydrogen atom or a $-\text{CO}(\text{C}_{1-4} \text{ alkyl})$, C_{1-6} alkyl, allyl, C_{3-6} cycloalkyl, phenyl or benzyl group;

R^{22} is a hydrogen atom or a C_{1-6} alkyl group;


R^{23} is a hydrogen atom or a C_{1-5} alkyl, aryl or $-\text{CH}_2\text{-aryl}$ group;

R^{24} is a hydrogen atom or a C_{1-4} alkyl group;

R^{25} is a hydrogen atom or a C_{1-8} alkyl, C_{1-8} perfluoroalkyl, C_{3-6} cycloalkyl, phenyl or benzyl group;

R^{26} is a hydrogen atom or a C_{1-6} alkyl, C_{3-6} cycloalkyl, phenyl or benzyl group;

R^{27} is $-\text{NR}^{23}\text{R}^{24}$, $-\text{OR}^{24}$, $-\text{NHCONH}_2$, $-\text{NHCSNH}_2$,



and

R^{28} and R^{29} are, independently of one another, each a C_{1-4} alkyl group or, taken together, a $-(CH_2)_q-$ group, where q is 2 or 3;

5 Wherein, R^{21} through R^{29} are with or without one or more substituents; and

(c) R^4 is a C_{3-15} cycloalkyl group, with or without substituents, such as a hydroxy substituent group, a C_{3-15} cycloalkenyl group, with or without one or more substituents, or a heterocycloalkyl group of 3 to 15 members, with or without one or more substituents;

10

wherein, the optional one or more substituents for all the groups are chemically-compatible and are, independently of one another, each defined the same as recited above in the definition section.

15

The invention comprises at least one compound of the formula (I), which includes any and all enantiomers, stereoisomers, rotomers, tautomers and prodrugs of the at least one inventive compound. Compounds of the formula (I) also include their corresponding salts, solvates (e.g., hydrates), esters, and the like. The invention further comprises pharmaceutically-acceptable compositions prepared from an inventive compound or a mixture of inventive compounds, or a salt, solvate or ester thereof. The compounds of formula (I) can be useful for treating a variety of diseases, symptoms and

20

physiological disorders, such as sexual dysfunction, especially impotence (e.g., erectile dysfunction).

A further understanding of the invention will be had from the following description of preferred embodiments.

5

DESCRIPTION OF PREFERRED EMBODIMENTS

The inventive compounds having the formula (I) are substituted at the 8-position on the chemical structure with an amino group that itself is substituted with one of the following groups: an unsaturated or saturated carbocyclic group and a saturated heterocyclic group. The inventive substituted xanthines exhibited unexpectedly enhanced properties with respect to enzyme activity and enzyme selectivity. It is believed that the substitution at the 8-position of the subject PDE V inhibitor compounds with these specific groups, helped produce unexpectedly highly potent and selective xanthines, which exhibited increased isozyme selectivity when compared to conventional xanthines. Pharmaceutical compositions comprising the inventive compounds possess unexpectedly superior therapeutic properties.

Referring above to the inventive xanthine PDE V inhibitor compounds having the formula (I), the 8-position on the chemical structure is substituted with a -NHR^4 group, where R^4 represents a carbocyclic or heterocyclic system defined as follows: a C_{3-15} cycloalkyl group, a C_{3-15} cycloalkenyl group or a heterocycloalkyl group of 3 to 15 members. All of the cyclic systems are optionally substituted. Preferred substituents on the cyclic systems include a

C₃₋₆ cycloalkyl group, a C₁₋₆ alkoxy C₁₋₆ alkyl group, a C₁₋₆ alkyl group, an amino C₁₋₆ alkyl group, a C₁₋₆ dialkylamino C₁₋₆ alkyl group, a C₃₋₆ dicycloalkylamino C₁₋₆ alkyl group, a hydroxy group, an alkoxy group, an oximino group, -COR⁶, -SO₂R⁶, -COOR⁶, -CONR⁶R⁷, -SO₂NR⁶R⁷,
 5 -N(R⁸)SO₂R⁶ and -NR⁶R⁷, where:


R⁶ is a hydrogen atom or an optionally substituted, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, aryl or heteroaryl group;

R⁷ is a hydrogen atom or an optionally substituted, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, aryl or heteroaryl group; or

10 R⁶ and R⁷, when applicable, may be joined together to form a heterocyclic ring system; and

R⁸ is a hydrogen atom or an optionally substituted, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, aryl or heteroaryl group.


Furthermore, R⁴ may also be substituted with -ZR⁷⁰Z'-, where R⁷⁰,
 15 together with Z and Z', form a spiro-fused 5- to 7-membered ring or a linearly fused 4- to 7-membered ring system, and Z and Z', independently of one another, are each an oxygen, sulfur or nitrogen atom. For example, when Z = Z' = O, R⁴ may be substituted by the following structure having the formula (VIII):



(VIII)

Preferred substituents are defined above for the groups. Other substituents may also be used, such as ketones, oximes, cyclic systems, including linearly fused and bridged, mono-, bi- and tricyclic rings, spiro-cyclic systems, including ketals and thioketals directly attached to R⁴, halogens and sulfonamides. One skilled in the art can determine other possible substituents depending on the conditions employed and the desired properties.

A preferred structure of the invention is represented by formula (II):



(II)

where,

R¹, R² and R³ are defined the same as above for the compound of formula (I);

R⁹ is one of the following atoms or groups:

- (a) a hydrogen atom;
- (b) an oximino group;
- (c) a carboxyalkyl group;
- (d) a C₁₋₆ alkoxy C₁₋₆ alkyl group;
- (e) an aryloxy C₁₋₆ alkyl group;
- (f) a C₃₋₆ cycloalkoxy C₁₋₆ alkyl group;

- (g) a heteroaryloxy C₁₋₆ alkyl group;
- (h) a -COOH group;
- (i) an ester group;
- (j) a C₁₋₆ alkyl group;
- (k) a C₃₋₆ cycloalkyl group;
- (l) a C₃₋₆ heterocyclic group;
- (m) a hydroxy C₁₋₆ alkyl group;
- (n) an aryl group; or
- (o) a heteroaryl group;

5

10

wherein, all of the above groups are optionally substituted;

R¹⁰ and R¹¹ are substituents on the same or different carbon atoms of the ring and, independently of one another, are each defined the same as above for R⁹ and, additionally, may each be one of the following groups:

15

- (a) a hydroxy group;
- (b) an ester group derived from a hydroxy group with
 - a:
 - (i) C₁₋₆ carboxylic acid;
 - (ii) C₃₋₆ cycloalkyl C₁₋₆ carboxylic acid;
 - (iii) aryl C₁₋₆ carboxylic acid; or
 - (iv) heteroaryl C₁₋₆ carboxylic acid group;
- (c) a C₁₋₆ alkoxy group;
- (d) an amino group;
- (e) a C₁₋₆ mono- or dialkylamino group;

20

- (f) a C₁₋₆ alkylacylamino group;
- (g) a C₁₋₆ alkylsulfonylamino group; or
- (h) a -NHCON(R¹⁴)₂ group, where R¹⁴ is a hydrogen atom or an optionally substituted, alkyl or aryl group; or

5 R¹⁰ and R¹¹, taken together with each other and, optionally, with one or more carbon and/or hetero atoms of the ring, form an optionally substituted, spiro-fused, linearly fused, bi- or tri-cyclic ring system of from 8 to 12 members, including from 0 to 4 hetero atoms, where, all of the above R¹⁰, R¹¹ and R¹⁴ groups are optionally substituted;

10 m and n are, independently of one another, each from 1 to 3; and

X is a chemcially-compatible group, which is -C(R¹⁰R¹¹)-, -S(O)_y, -O-, -N(R⁶⁰)-, where:

15 R¹⁰ and R¹¹ are, independently of one another, each defined the same as previously;

y is from 0 to 2;

15 R⁶⁰ is a hydrogen atom or a C₁₋₈ alkyl, C₁₋₈ alkynyl, C₁₋₈ alkenyl, C₃₋₈ cycloalkyl, aryl, heteroaryl, C₄₋₈ heterocycloalkyl, COR⁶¹, SO₂R⁶¹, COOR⁶¹, CONR⁶¹R⁶² or SO₂NR⁶¹R⁶² group, with or without substituents, where:

20 R⁶¹ is a hydrogen atom or a C₁₋₈ alkyl, C₁₋₈ alkynyl, C₁₋₈ alkenyl, C₃₋₈ cycloalkyl, aryl, heteroaryl or C₄₋₈ heterocyclic group, with or without substituents;

R^{62} is a hydrogen atom or a C_{1-8} alkyl, C_{1-8} alkynyl, C_{1-8} alkenyl, C_{3-8} cycloalkyl, aryl, heteroaryl or C_{4-8} heterocyclic group, with or without substituents; and when R^{61} and R^{62} are (the same or different) alkyl groups, they can, if desired, be joined together to form a carbocyclic or heterocyclic ring system;

wherein, the optional substituents and the one or more substituents are defined the same as for the one or more substituents of formula (I) above.

In the compound of formula (II), the different carbon atoms to which R^{10} and R^{11} may be connected can be adjacent or non-adjacent. Preferably, R^9 , R^{10} and R^{11} are all hydrogen atoms. In another embodiment of the invention, one of R^{10} or R^{11} is, advantageously, a hydroxy group.

In the compounds of formulas (I) and (II), R^1 is, preferably, an alkyl group or an arylalkyl group, particularly, a benzyl group. More preferably, R^1 is a lower alkyl group of from 1 to 4 carbon atoms, and most preferably, a methyl or ethyl group.

R^2 , in the compounds of formulas (I) and (II), is, preferably, an alkyl group, particularly, an alkyl group substituted with a hydroxy group. More preferably, R^2 is a lower alkyl group of from 1 to 3 carbon atoms or a hydroxyalkyl group, and most preferably, R^2 is a methyl, ethyl, iso-butyl or hydroxyethyl group.

In the compounds of formulas (I) and (II), R^3 is, preferably, an aryl group, particularly, an aryl group substituted with a hydroxy-, alkoxy- or

amino-sulfonyl group, which may be, advantageously, substituted with 1 or 2 halogen atoms. When R^3 is a heteroaryl group in the compounds of formulas (I) and (II), it is generally preferable to utilize heteroaryl groups other than furan. Most preferably, R^3 is a methoxyaryl group substituted on its aryl ring with at least one halogen atom, for example, a substitution with 1 or 2 halogen atoms, such as chlorine or bromine. For instance, R^3 can be 4-hydroxyphenyl, 3-chloro-4-hydroxyphenyl, 3-bromo-4-hydroxyphenyl, 4-methoxyphenyl, 3-chloro-4-methoxyphenyl, 3-bromo-4-methoxyphenyl, 4-aminosulfonylphenyl group, 3-chloro-4-aminosulfonylphenyl group or 3-bromo-4-aminosulfonyl-phenyl.

R^4 , in the compound of formula (I), is, preferably, a cycloalkyl or heterocycloalkyl group, particularly, a cycloalkyl group substituted with a hydroxy group. More preferably, R^4 is a cyclohexyl, hydroxycyclopentyl or tetrahydropyranyl group. Most preferably, R^4 is a hydroxycyclopentyl group. For instance, R^4 can be a 2(R)-hydroxy-1(R)-cyclopentyl group. All of the preferred embodiments may be unsubstituted or substituted.

The compounds of formulas (I) and (II) are useful for treating urogenital diseases, such as male (e.g., impotence / erectile dysfunction) and female sexual dysfunction. The following compounds listed in Tables I and II are illustrative of the invention:








TABLE I	
Compound No.	Structure
10	
11	
12	
13	
14	
15	
16	







TABLE I	
Compound No.	Structure
17	
18	
19	
20	
21	
22	

TABLE I	
Compound No.	Structure
23	
24	
25	
28	
29	
30	
31	
32	









TABLE I	
Compound No.	Structure
33	
34	
35	
36	
37	
38	
39	
40	








TABLE I	
Compound No.	Structure
41	
42	
43	
44	
47	
48	
49	








TABLE I	
Compound No.	Structure
50	
51	
52	
53	
54	
55	
56	








TABLE I	
Compound No.	Structure
57	
58	
59	
60	
61	
62	
63	








TABLE I	
Compound No.	Structure
64	
65	
66	
67	
68	
69	
70	









TABLE I	
Compound No.	Structure
71	
72	
73	
74	
75	
76	
77	
78	

TABLE I	
Compound No.	Structure
79	
80	
81	
82	
83	
84	
85	

TABLE I	
Compound No.	Structure
86	
87	
88	
89	
90	
91	
92	
93	







TABLE I	
Compound No.	Structure
94	
95	
96	
97	
98	
99	

TABLE II







Compound No.	STRUCTURE	HRMS Calc.	HRMS Found	M, M+1
100		439.2821	439.2821	(M+1)
101		412.2349	412.2346	(M+1)
102		526.3213	526.3203	(M+)
103		442.2454	442.2451	(M+1)
104		428.2298	428.2294	(M+1)
105		476.2065	476.2057	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
106		478.1857	478.1851	(M+1)
107		462.1908	462.191	(M+1)
108		490.1857	490.1853	(M+1)
109		492.1650	492.1641	(M+1)
110		455.2533	455.2518	(M+)
111		458.2403	458.2395	(M+1)

TABLE II







Compound No.	STRUCTURE	HRMS Calc.	HRMS Found	M, M+1
112		442.2454	442.2448	(M+1)
113		444.2247	444.2252	(M+1)
114		522.1352	522.1346	(M+1)
115		464.1701	464.1696	(M+1)
116		506.1403	506.141	(M+1)
117		520.1559	520.1568	(M+1)







TABLE II				
<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
118		508.1196	508.119	(M+1)
119		475.2128	475.2134	(M+1)
120		429.1932	429.1931	(M+)
121		488.2332	488.2333	(M+1)
122		504.1610	504.1605	(M+1)
123		506.1403	506.1395	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
124		522.1542	522.1542	(M+1)
125		520.1559	520.1552	(M+1)
126		477.1920	477.1919	(M+1)
127		477.1920	477.1914	(M+1)
128		536.1335	536.1335	(M+1)
129		522.1352	522.136	(M+1)

TABLE II






<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
130	no structure	n/a	n/a	n/a
131		382.2243	382.2242	(M+1)
132		382.2243	382.2238	(M+1)
133		424.2713	424.2717	(M+1)
134		396.2400	396.2396	(M+1)
135		396.2400	396.2393	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
136		386.1992	386.1988	(M+1)
137		386.1992	386.1988	(M+1)
138		386.1992	386.1985	(M+1)
139		398.2192	398.2196	(M+1)
140		382.2243	382.2238	(M+1)
141		398.2192	398.2192	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
142		412.1985	412.1982	(M+1)
143		428.2298	428.2294	(M+1)
144		412.2349	412.2346	(M+1)
145		384.2036	384.2041	(M+1)
146		384.2036	384.2033	(M+1)
147		398.2192	398.2184	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
148		402.1697	402.1691	(M+1)
149		493.0975	493.098	(M+)
150		451.1831	451.1819	(M+)
151		435.1882	435.1879	(M+)
152		446.1192	446.1187	(M+1)
153		435.1229	435.1219	(M+)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
154		404.1898	404.1895	(M+1)
155		428.2298	428.2292	(M+1)
156		420.1603	420.1603	(M+1)
157		413.1937	413.1932	(M+1)
158		444.2400	444.2394	(M+1)
159		431.1724	431.173	(M+)







TABLE II				
<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
160		446.1595	446.1588	(M+1)
161		418.1646	418.164	(M+1)
162		436.1960	436.1962	(M+1)
163		436.1960	436.1957	(M+1)
164		452.1909	452.1919	(M+1)
165		414.2305	414.2303	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
166		440.2662	440.2657	(M+1)
167		426.2505	426.2509	(M+1)
168		440.2298	440.2295	(M+1)
169		426.2505	426.2498	(M+1)
170		412.2349	412.2345	(M+1)
171		474.2272	474.2277	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
172		459.2037	459.2055	(M+)
173		428.2462	428.2457	(M+1)
174		440.2662	440.2657	(M+1)
175		454.2454	454.2449	(M+1)
176		454.2818	454.2812	(M+1)
177		426.2505	426.2503	(M+1)

TABLE II







<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
178		440.2662	440.2666	(M+1)
179		509.1738	509.1729	(M+1)
180		555.1233	555.123	(M+1)
181		511.153	511.1524	(M+1)
182		491.2077	491.2087	(M+1)
183		525.1687	525.1697	(M+1)

TABLE II







Compound No.	STRUCTURE	HRMS Calc.	HRMS Found	M, M+1
184		571.1164	571.1138	(M+1)
185		538.1492	538.1498	(M+1)
186		524.1335	524.1344	(M+1)
187		575	575	(M) LRMS
188		477.192	477.1919	(M+1)
189		477.192	477.1919	(M+1)

TABLE II











<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M. M+1</u>
190		557.1007	557.0997	(M+1)
191		511.153	511.1519	(M+1)
192		494.1637	494.1636	(M+1)
193		510.1578	510.1574	(M+1)
194		554.1073	554.1066	(M+1)
195		525.159	525.1582	(M+1)

TABLE II				
<u>Compound No.</u>	<u>STRUCTURE</u>	<u>HRMS Calc.</u>	<u>HRMS Found</u>	<u>M, M+1</u>
196		525.159	525.1597	(M+1)
197		478.168	478.1683	(M+1)
198		522.1174	522.1169	(M+1)
199		542.1405	542.143	(M+1)

The compounds of the invention are useful for inhibiting PDE V enzymes. Their enzyme activities and enzyme selectivities can be evaluated in a number of ways. In particular, enzyme activity can be measured by the PDE V IC₅₀ value, which is the concentration (in nM) of the compound required to provide 50% inhibition of PDE V. The lower the value of IC₅₀, the

more active is the compound. Measurements on the compounds in Tables I and II gave the following data (all numbers are modified by the word "about"):

- 5 A. all compounds had a PDE V IC_{50} within the range of from < 1 nM to >100 nM;
- B. compound nos. 13-18, 25, 30-32, 38, 41-43, 55-58, 69-71, 77, 85, 92, 96, 98, 101, 113, 120, 121, 126, 128, 131, 137, 138, 141, 146-48, 165, 166, 173, 176, 181, 182, 184, 185, 193 and 194 had a PDE V IC_{50} within the range of from > 15 to 100 nM;
- 10 C. compound nos. 23, 24, 29, 33, 34, 39, 40, 93, 94, 108, 111, 112, 125, 136, 144, 160 and 161 had a PDE V IC_{50} within the range of from > 10 to 15 nM.
- D. compound nos. 21, 22, 28, 36, 37, 59, 66, 68, 78, 79, 89, 95, 99, 110, 115, 132, 159, 171, 172, 175, 180, 183, 190 and 199 had a
- 15 PDE V IC_{50} within the range of from > 5 to 10 nM; and
- E. compound nos. 60-65, 67, 103-07, 114, 116-19, 122-24, 142, 168-70, 177, 178, 179, 186-88, 191, 197 and 198 had a PDE V IC_{50} within the range of up to 5 nM.


20 In addition, another type of measurement that can be made is the ratio of PDE VI IC_{50} / PDE V IC_{50} (identified as "PDE VI / PDE V"), which is an indicator of enzyme selectivity – the higher the ratio, the more selective is the compound to inhibiting PDE V enzyme relative to PDE VI enzyme. Measurements on the compounds (except for compound nos. 189, 192, 195

and 196) in Table II gave the following data (all numbers are modified by the word "about"):



- 5 F. compound nos. 1-188, 190, 191, 193, 194 and 197-99 had a
PDE VI / PDE V ratio of > 0 ;
- G. compound nos. 165 and 193 had a PDE VI / PDE V ratio within
the range of from > 0 to 10;
- H. compound nos. 101, 108, 136, 141, 146, 148, 168, 173 and 194
had a PDE VI / PDE V ratio within the range of from > 10 to 25;
- 10 I. compound nos. 104, 125, 131-32, 137-38, 142, 144, 170, 175,
177, 185 and 199 had a PDE VI / PDE V ratio within the range
of from > 25 to 50;
- J. compound nos. 103, 110, 111, 117, 159, 166, 182 and 187 had
a PDE VI / PDE V ratio within the range of from > 50 to 75;
- 15 K. compound nos. 105, 106, 147 and 171 had a PDE VI / PDE V
ratio within the range of from > 75 to 100;
- L. compound nos. 112, 113, 123, 124, 126, 169, 172 and 184 had
a PDE VI / PDE V ratio within the range of from > 100 to 140;
and
- 20 M. compound nos. 107, 114-16, 118-22, 128, 160-61, 176, 178-81,
183, 186, 188, 190, 191, 197 and 198 had a PDE VI / PDE V
ratio of from > 140 .

Preferred compounds of the invention include those found in classes E and/or M: compound nos. 60-65, 67, 103-07, 114-24, 128, 142, 160-61, 168-70, 176-78, 179, 186, 188, 191, 197 and 198. More preferred compounds of the invention are compound nos. 107, 114, 116, 118, 119, 122, 160, 178 and 186 of Table II.

Another preferred compound of the invention would have the following chemical structure:



Specific and general procedures for producing three preferred compounds follow below (compound nos. 107, 114 and 160). Obvious modifications to these procedures may be undertaken by one of ordinary skill in the art. Other compounds of the invention may be produced along the same lines.

Synthesis of Compound No. 107 in Table II (7)Preparation of 9

Experimental Procedure: Compound No. 107 in Table II (7)

1 (20.0 g, 74.0 mmol) was dissolved in dimethylformamide (370 mL) under nitrogen and (2-bromoethoxy)-*tert*-butyldimethylsilane (31.8 mL, 148 mmol) was added dropwise. The reaction was stirred at room temperature for 115 hrs., then diluted with ethyl acetate and washed with water several times. The organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (30/70 ethyl acetate/hexanes) yielded **2** (28.1 g, 88%).

^1H NMR (400 MHz, CDCl_3): δ 7.52 (s, 1H), 7.29-7.39 (m, 5H), 5.49 (s, 2H), 4.25 (t, 2H, $J = 6.0$ Hz), 4.07 (q, 2H, $J = 7.2$ Hz), 3.93 (t, 2H, $J = 6.0$ Hz), 1.24 (t, 3H, $J = 7.2$ Hz), 0.75 (s, 9H), 0.08 (s, 6H). HRMS: Calcd for $\text{C}_{22}\text{H}_{32}\text{N}_4\text{O}_3\text{Si}$ (M+H): 429.2322. Found: 429.2329.

To a solution of **2** (2.10 g, 4.89 mmol) in methanol (375 mL) was added ammonium formate (4.64g, 73.6 mmol) and 20% palladium hydroxide on carbon (980 mg). The reaction was heated to reflux for 1.5 hrs., then cooled to room temperature, filtered and concentrated under vacuum. Purification via flash chromatography (50/50 ethyl acetate/hexanes) yielded **3** (1.26 g, 94%).

^1H NMR (400 MHz, CDCl_3): δ 7.82 (s, 1H), 4.33 (t, 2H, $J = 6.0$ Hz), 4.16 (q, 2H, $J = 7.2$ Hz), 3.99 (t, 2H, $J = 6.0$ Hz), 1.29 (t, 3H, $J = 7.2$ Hz), 0.78 (s, 9H), 0.06 (s, 6H). HRMS: Calcd for $\text{C}_{15}\text{H}_{26}\text{N}_4\text{O}_3\text{Si}$ (M+H): 339.1852. Found: 339.1864.

To **3** (970 mg, 2.86 mmol) was added dimethylformamide (14 mL), 3-chloro-4-methoxybenzyl bromide **9** (1.72 g, 5.70 mmol), and potassium carbonate (785 mg, 5.70 mmol) under nitrogen. The reaction mixture was stirred at room temperature for 24 hrs., then diluted with ethyl acetate and washed with water several times. The organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification by flash chromatography (30/70 ethyl acetate/hexanes) yielded **4** (1.14 g, 81%).

¹H NMR (400 MHz, CDCl₃): δ 7.52 (s, 1H), 7.33 (d, 1H, *J* = 2.4 Hz), 7.25 (dd 1H, *J* = 2.0 Hz, *J* = 8.4 Hz), 6.90 (d, 1H, *J* = 8.8 Hz), 5.40 (s, 2H), 4.25 (t, 2H, *J* = 6.0 Hz), 4.07 (q, 2H, *J* = 7.2 Hz), 3.93 (t, 2H, *J* = 6.0 Hz), 3.89 (s, 3H), 1.25 (t, 3H, *J* = 7.2 Hz), 0.75 (s, 9H), 0.08 (s, 6H). HRMS: Calcd for C₂₃H₃₃ClN₄O₄Si (M+H): 493.2038. Found: 493.2032.

To solution of **4** (1.14g, 2.32 mmol) in tetrahydrofuran (20 mL) under nitrogen at -78 °C (dry ice/acetone bath) was added lithium diisopropylamide (2M in THF/heptane, 1.7 mL, 3.48 mmol). After stirring for thirty minutes, 1,2-dibromotetrafluoroethane (0.55 mL, 4.63 mmol) was added dropwise over five minutes. The reaction was stirred for 1.5 hrs. at -78 °C then quenched with saturated aqueous sodium bicarbonate and warmed to room temperature. The mixture was extracted with dichloromethane, dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (30/70 ethyl acetate/hexanes) yielded **5** (640 mg, 48%).

¹H NMR (400 MHz, CDCl₃): δ 7.42 (d, 1H, *J* = 2.4 Hz), 7.31 (dd, 1H, *J* = 2.0 Hz, *J* = 8.4 Hz), 6.88 (d, 1H, *J* = 8.8 Hz), 5.45 (s, 2H), 4.22 (t, 2H, *J*

= 5.6 Hz), 4.07 (q, 2H, J = 7.2 Hz), 3.92 (t, 2H, J = 5.6 Hz), 3.88 (s, 3H), 1.25 (t, 3H, J = 7.2 Hz), 0.74 (s, 9H), 0.08 (s, 6H). HRMS: Calcd for $C_{23}H_{32}BrClN_4O_4Si$ (M+H): 571.1143. Found: 571.1149.

5 To **5** (320 mg, 0.56 mmol) was added cyclohexylamine (0.25 mL, 2.24 mmol), diisopropylethylamine (2.8 mL), and 1-methyl-2-pyrrolidinone (2.8 mL). The reaction mixture was heated to 160 °C in a sealed tube for 18 hrs., then cooled to room temperature. Water was added, then the mixture was extracted with ethyl acetate and washed with water several times. The
10 organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (30/70 ethyl acetate/hexanes) yielded **6** (210 mg, 64%).

1H NMR (400 MHz, $CDCl_3$): δ 7.27 (d, 1H, J = 2.0 Hz), 7.13 (dd, 1H, J = 2.0 Hz, J = 8.4 Hz), 6.89 (d, 1H, J = 8.8 Hz), 5.23 (s, 2H), 4.19 (t, 2H, J = 6.2 Hz), 4.05 (q, 2H, J = 7.2 Hz), 3.93 (t, 2H, J = 6.2 Hz), 3.89 (s, 3H), 3.86-3.91 (m, 1H), 3.69-3.80 (m, 1H), 1.88-1.96 (m, 2H), 1.52-1.64 (m, 3H), 1.28-1.42 (m, 2H), 1.23 (t, 3H, J = 7.2 Hz), 1.04-1.22 (m, 3H), 0.81 (s, 9H), 0.01 (s, 6H). HRMS: Calcd for $C_{29}H_{44}ClN_5O_4Si$ (M+H): 590.2937. Found: 590.2929.

20 **6** (191 mg, 0.324 mmol) was dissolved in dichloromethane (4.0 mL) under nitrogen and cooled to 0 °C in an ice bath. Boron tribromide (0.14 mL, 1.42 mmol) was added to the reaction mixture and warmed to room temperature. After 1.25 hr., the reaction was diluted with dichloromethane

and washed with water several times. The organic phase was dried over potassium carbonate, filtered and concentrated under vacuum. Purification via PTLC (70/30 ethyl acetate/hexanes) yielded **7** (compound no. 107 in Table II) (122 mg, 74%).

5 ¹H NMR (400 MHz, CDCl₃): δ 7.25-7.28 (m, 1H), 7.09 (dd, 1H, *J* = 2.0 Hz, *J* = 8.0 Hz), 7.01 (d, 1H, *J* = 8.0 Hz), 5.69 (s, 1H), 5.23 (s, 2H), 4.32-4.36 (m, 2H), 4.16 (t, 1H, *J* = 6.0 Hz), 4.06 (q, 2H, *J* = 7.2 Hz), 3.90-3.98 (m, 3H), 3.62-3.72 (m, 1H), 1.87-1.96 (m, 2H), 1.54-1.66 (m, 3H), 1.31-1.43 (m, 2H), 1.25 (t, 3H, *J* = 7.2 Hz), 1.06-1.22 (m, 3H).

10 HRMS: Calcd for C₂₂H₂₈ClN₅O₄ (M+H): 462.1908. Found: 462.1901.


3-Chloro-4-methoxytoluene **8** (2.6 mL, 19.2 mmol) was dissolved in dichloromethane (30 mL) and *N*-bromosuccinimide (3.75 g, 21.1 mmol) was added followed by AIBN (36.0 mg). The reaction was heated to reflux for 19
15 hrs., then cooled to room temperature and the precipitate was filtered off. The filtrate was diluted with dichloromethane and washed with 0.5 M aqueous sodium bicarbonate, followed by water. The organic mixture was dried over sodium sulfate, filtered and concentrated under vacuum to yield **9** (4.73g, 82%). The benzyl bromide was used as the crude material without further
20 purification.

¹H NMR (400 MHz, CDCl₃): δ 7.42 (d, 1H, *J* = 2.4 Hz), 7.26 (dd, 1H, *J* = 2.4 Hz, *J* = 8.4 Hz), 6.88 (d, 1H, *J* = 8.4 Hz), 4.44 (s, 2H), 3.90 (s, 3H).


General Synthesis of Compound No. 107 in Table II (7)

- a) Reacting **1** with an alkyl halide and base to form **2**;
- b) Debenzylation of **2** to form **3**;
- c) Alkylation of **3** with a benzyl halide to form **4**;
- 5 d) Deprotonation of **4** followed by addition of a brominating agent to form **5**;
- e) Displacement of bromo **5** with an amine to form **6**; and
- f) Treatment of **6** with boron tribromide to form compound no. 107 in Table II (7) via cleavage of both silyl and methyl ethers.

Synthesis of Compound 114 in Table II (13)



Preparation of 15



Experimental Procedure: Compound 114 in Table II (13)

1 (20.0 g, 74.0 mmol) was dissolved in dimethylformamide (370 mL) under nitrogen and (2-bromoethoxy)-*tert*-butyldimethylsilane (31.8 mL, 148 mmol) was added dropwise. The reaction was stirred at room temperature for 115 hrs., then diluted with ethyl acetate and washed with water several times. The organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (30/70 ethyl acetate/hexanes) yielded **2** (28.1 g, 88%).

¹H NMR (400 MHz, CDCl₃): δ 7.52 (s, 1H), 7.29-7.39 (m, 5H), 5.49 (s, 2H), 4.25 (t, 2H, *J* = 6.0 Hz), 4.07 (q, 2H, *J* = 7.2 Hz), 3.93 (t, 2H, *J* = 6.0 Hz), 1.24 (t, 3H, *J* = 7.2 Hz), 0.75 (s, 9H), 0.08 (s, 6H). HRMS: Calcd for C₂₂H₃₂N₄O₃Si (M+H): 429.2322. Found: 429.2329.

To a solution of **2** (2.10 g, 4.89 mmol) in methanol (375 mL) was added ammonium formate (4.64g, 73.6 mmol) and 20% palladium hydroxide on carbon (980 mg). The reaction was heated to reflux for 1.5 hrs., then cooled to room temperature, filtered and concentrated under vacuum. Purification via flash chromatography (50/50 ethyl acetate/hexanes) yielded **3** (1.26 g, 94%).

¹H NMR (400 MHz, CDCl₃): δ 7.82 (s, 1H), 4.33 (t, 2H, *J* = 6.0 Hz), 4.16 (q, 2H, *J* = 7.2 Hz), 3.99 (t, 2H, *J* = 6.0 Hz), 1.29 (t, 3H, *J* = 7.2 Hz), 0.78 (s, 9H), 0.06 (s, 6H). HRMS: Calcd for C₁₅H₂₆N₄O₃Si (M+H): 339.1852. Found: 339.1864.

To **3** (970 mg, 2.86 mmol) was added dimethylformamide (25 mL), 3-bromo-4-methoxybenzyl bromide **15** (1.62 g, 5.79 mmol), and potassium carbonate (800 mg, 5.79 mmol) under nitrogen. The reaction mixture was stirred at room temperature for 21 hrs., then diluted with ethyl acetate and washed with water several times. The organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification by flash chromatography (30/70 ethyl acetate/hexanes) yielded **10** (1.55 g, 100%).

¹H NMR (400 MHz, CDCl₃): δ 7.52 (s, 1H), 7.51 (d, 1H, *J* = 2.4 Hz), 7.30 (dd 1H, *J* = 2.0 Hz, *J* = 8.4 Hz), 6.87 (d, 1H, *J* = 8.8 Hz), 5.40 (s, 2H), 4.25 (t, 2H, *J* = 6.0 Hz), 4.07 (q, 2H, *J* = 7.0 Hz), 3.93 (t, 2H, *J* = 6.0 Hz), 3.88 (s, 3H), 1.25 (t, 3H, *J* = 7.0 Hz), 0.75 (s, 9H), 0.08 (s, 6H).
HRMS: Calcd for C₂₃H₃₃BrN₄O₄Si (M+H): 537.1533. Found: 537.1540.

To solution of **10** (1.50 g, 2.80 mmol) in tetrahydrofuran (24 mL) under nitrogen at -78 °C (dry ice/acetone bath) was added lithium diisopropylamide (2M in THF/heptane, 2.2 mL, 4.33 mmol). After stirring for thirty minutes, 1,2-dibromotetrafluoroethane (0.69 mL, 5.77 mmol) was added dropwise over five minutes. The reaction was stirred for 1.25 hrs. at -78 °C then quenched with saturated aqueous sodium bicarbonate and warmed to room temperature. The mixture was extracted with dichloromethane, dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (30/70 ethyl acetate/hexanes) yielded **11** (600 mg, 34%).

¹H NMR (400 MHz, CDCl₃): δ 7.60 (d, 1H, *J* = 2.4 Hz), 7.35 (dd, 1H, *J* = 2.0 Hz, *J* = 8.4 Hz), 6.84 (d, 1H, *J* = 8.4 Hz), 5.45 (s, 2H), 4.21 (t, 2H, *J* = 5.6 Hz), 4.07 (q, 2H, *J* = 6.8 Hz), 3.90 (t, 2H, *J* = 5.6 Hz), 3.87 (s, 3H), 1.24 (t, 3H, *J* = 6.8 Hz), 0.73 (s, 9H), 0.08 (s, 6H). HRMS: Calcd for C₂₃H₃₂Br₂N₄O₄Si (M+H): 615.0638. Found: 615.0633.

To **11** (1.89 g, 3.07 mmol) was added the amino alcohol hydrochloride salt (1.31 g, 12.27 mmol), diisopropylethylamine (15.4 mL), and 1-methyl-2-pyrrolidinone (15.4 mL). The reaction mixture was heated to 160 °C in a sealed tube for 13 hrs., then cooled to room temperature. Water was added, then the mixture was extracted with ethyl acetate and washed with water several times. The organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (3/97 methanol/dichloromethane) yielded **12** (1.77 g, 90%).

¹H NMR (400 MHz, CDCl₃): δ 7.45 (d, 1H, *J* = 2.0 Hz), 7.17 (dd, 1H, *J* = 2.4 Hz, *J* = 8.6 Hz), 6.86 (d, 1H, *J* = 8.4 Hz), 5.18-4.34 (m, 3H), 4.00-4.23 (m, 5H), 3.86-3.98 (m, 6H), 3.69-3.79 (m, 1H), 2.10-2.21 (m, 1H), 1.99-2.10 (m, 1H), 1.60-1.84 (m, 3H), 1.32-1.43 (m, 1H), 1.24 (t, 3H, *J* = 7.2 Hz), 0.75 (s, 9H), 0.07 (d, 6H, *J* = 4.0 Hz). HRMS: Calcd for C₂₈H₄₃BrN₅O₅Si (M+H): 636.2217. Found: 636.2207.

12 (1.77 g, 2.78 mmol) was dissolved in tetrahydrofuran (28 mL) under nitrogen and tetrabutylammonium fluoride (1M in THF, 28 mL) was added dropwise. The reaction was stirred at room temperature for 15 hrs., then

diluted with dichloromethane and washed with water several times. The organic mixture was dried over potassium carbonate, filtered and concentrated under vacuum. Purification via flash chromatography (3/97 methanol/dichloromethane) yielded **13** (compound no. 114 in Table II) (760 mg, 52%).

¹H NMR (400 MHz, CDCl₃): δ 7.47 (d, 1H, *J* = 2.0 Hz), 7.19 (dd, 1H, *J* = 2.0 Hz, *J* = 8.4 Hz), 6.88 (d, 1H, *J* = 8.4 Hz), 5.25 (s, 2H), 5.09 (s, 1H), 4.21-4.27 (m, 3H), 4.06 (q, 2H, *J* = 7.0 Hz), 3.90-3.97 (m, 3H), 3.89 (s, 1H), 3.74-3.82 (m, 1H), 3.08 (s, 1H), 2.12-2.22 (m, 1H), 1.98-2.08 (m, 1H), 1.60-1.86 (m, 3H), 1.33-1.43 (m, 1H), 1.25 (t, 3H, *J* = 7.0 Hz), 1.06-1.22 (m, 3H). HRMS: Calcd for C₂₂H₂₈BrN₅O₅ (M+H): 522.1352. Found: 522.1346.


2-Bromo-4-methyl anisole **14** (2.2 mL, 14.9 mmol) was dissolved in dichloromethane (30 mL) and N-bromosuccinimide (3.75 g, 16.4 mmol) was added followed by AIBN (26.0 mg). The reaction was heated to reflux for 19 hrs., then cooled to room temperature and the precipitate was filtered off. The filtrate was diluted with dichloromethane and washed with 0.5 M aqueous sodium bicarbonate, followed by water. The organic mixture was dried over sodium sulfate, filtered and concentrated under vacuum to yield **15** (4.16 g, 100%). The benzyl bromide was used as the crude material without further purification.

¹H NMR (400 MHz, CDCl₃): δ 7.59 (d, 1H, *J* = 2.0 Hz), 7.30 (dd, 1H, *J* = 2.4 Hz, *J* = 8.4 Hz), 6.85 (d, 1H, *J* = 8.4 Hz), 4.37 (s, 2H), 3.90 (s, 3H).

General Synthesis of Compound No. 114 in Table II (13)

- a) Reacting **1** with an alkyl halide and base to form **2**;
b) Debenzylation of **2** to form **3**;
c) Alkylation of **3** with a benzyl halide to form **10**;
d) Deprotonation of **10** followed by addition of a brominating agent to form **11**;
e) Displacement of bromo **11** with an amine to form **12**; and
f) Silyl ether cleavage of **12** to form compound no. 114 in Table II (13).

Synthesis of Compound No. 160 in Table II (18)



Experimental Procedure: Compound No. 160 in Table II (18)

To **16** (150 mg, 0.579 mmol) was added dimethylformamide (3 mL), 6-chloropiperonyl chloride (142 mg, 0.694 mmol) and potassium carbonate (120

mg, 0.868 mmol) under nitrogen. The reaction mixture was stirred at room temperature for 137 hrs., then diluted with ethyl acetate and washed with water several times. The organic mixture was dried over magnesium sulfate, filtered and concentrated under vacuum. Purification via PTLC (1/1 ethyl acetate/hexanes) yielded **17** (84.1 mg, 34%).

^1H NMR (400 MHz, CDCl_3): δ 6.89 (s, 1H), 6.09 (s, 1H), 5.95 (s, 2H), 5.59 (s, 2H), 3.60 (s, 3H), 3.38 (s, 3H). HRMS: Calcd for $\text{C}_{15}\text{H}_{12}\text{BrClN}_4\text{O}_4$ (M+H): 426.9809. Found: 426.9802.

To **17** (72.0 mg, 0.169 mmol) was added cyclohexylamine (86.7 mg, 0.883 mmol), diisopropylethylamine (0.8 mL) and 1-methyl-2-pyrrolidinone (0.8 mL). The reaction mixture was heated to 160 °C in a sealed tube for 17 hrs., then cooled to room temperature. Water was added, then the mixture was extracted with ethyl acetate and washed with water several times. The organic mixture was dried over magnesium sulfate, filtered and concentrated under vacuum. Purification via PTLC (1/1 ethyl acetate/hexanes) yielded compound no. 160 in Table II (**18**) (50.9 mg, 68%).

^1H NMR (400 MHz, CDCl_3): δ 6.84 (s, 1H), 6.815 (s, 1H), 5.96 (s, 2H), 5.33 (s, 2H), 4.42 (d, 1H, J = 7.2 Hz), 3.68-3.79 (m, 1H), 3.53 (s, 3H), 3.40 (s, 3H), 1.94-2.05 (m, 2H), 1.55-1.74 (m, 3H), 1.25-1.42 (m, 2H), 1.10-1.22 (m, 3H). HRMS: Calcd for $\text{C}_{21}\text{H}_{24}\text{ClN}_5\text{O}_4$ (M+H): 444.2400. Found: 444.2394.

General Synthesis of Compound No. 160 in Table II (18)

a) Alkylation of **16** with a benzyl halide to form **17**; and


b) Displacement of bromo **17** with an amine to form compound no. 160 in Table II (**18**).

5

Accordingly, the invention includes a method for producing a compound having the formula (I), comprising:

- (i) reacting a compound having the formula (III) with an alkyl halide in the presence of a base to form a compound having the formula (IV):

10



where,

15

- (a) R^1 is a hydrogen atom or a C_{1-15} alkyl group, branched or straight chain, with or without one or more substituents, a C_{2-15} alkenyl group, branched or straight chain, with or without one or more substituents, a C_{2-15} alkynyl group, branched or straight chain, with or without one or more substituents, a C_{3-15} cycloalkyl group, with or without one or more substituents, an arylalkyl group, with or without one or more substituents, an aryl group, with or without one or more substituents, a heteroaryl group, with or without one or more substituents, $-OR^5$, $-COOR^5$, $-C(O)R^5$ or $-C(O)N(R^5)_2$, where

20


R^5 is a hydrogen atom or a hydrocarbon radical, branched or straight-chain, with or without one or more substituents

(b) L is R^2 or a protected form of R^2 ; and

(c) Ph is a phenyl group;

5

(ii) debenzylating and then alkylating the compound having the formula (IV) with an alkyl halide, XCH_2R^3 , to form the compound having the formula (V):



where,

X is a halogen atom (e.g., a chlorine or bromine atom)


and

15

R^3 is an aryl group, with or without one or more substituents, a heteroaryl group, with or without one or more substituents, or a heterocyclic group having 1 to 3 heteroatoms fused to a 5 or 6 membered aryl ring, with or without one or more substituents, with the proviso that R^3 is not an aryl group substituted at its para position with a -Y-aryl group, where Y is a carbon-carbon single bond, -CO-, -O-, -S-, -N(R^{21})-, -CON(R^{22})-, -N(R^{22})CO-, -OCH₂-, -CH₂O-, -SCH₂-, -CH₂S-, -NHC(R^{23})(R^{24})-,

20

$-\text{NR}^{23}\text{SO}_2-$, $-\text{SO}_2\text{NR}^{23}-$, $-\text{C}(\text{R}^{23})(\text{R}^{24})\text{NH}-$, $-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$,
 $-\text{CH}=\text{CF}-$, $-\text{CF}=\text{CH}-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CF}_2\text{CF}_2-$,



5

where,

R^{21} is a hydrogen atom or a $-\text{CO}(\text{C}_{1-4}$ alkyl), C_{1-6} alkyl, allyl, C_{3-6} cycloalkyl, phenyl or benzyl group;

R^{22} is a hydrogen atom or a C_{1-6} alkyl group;

R^{23} is a hydrogen atom or a C_{1-5} alkyl, aryl or $-\text{CH}_2$ -aryl

10

group;

R^{24} is a hydrogen atom or a C_{1-4} alkyl group;

R^{25} is a hydrogen atom or a C_{1-8} alkyl, C_{1-8} perfluoroalkyl;


C_{3-6} cycloalkyl, phenyl or benzyl group;

R^{26} is a hydrogen atom or a C_{1-6} alkyl, C_{3-6} cycloalkyl, phenyl

15

or benzyl group;

R^{27} is $-\text{NR}^{23}\text{R}^{24}$, $-\text{OR}^{24}$, $-\text{NHCONH}_2$, $-\text{NHCSNH}_2$,



and

R^{28} and R^{29} are, independently of one another, each a C_{1-4} alkyl group, or R^{28} and R^{29} , taken together with each other, are a $-(CH_2)_q$ group, where q is 2 or 3;

wherein, R^{21} through R^{29} are optionally substituted with any of the groups defined above for the one or more substituents; and


(iii) deprotonating and then halogenating the compound having the formula (V) to form a compound having the formula (VI):



where,

Hal is a halogen atom;


(iv) reacting the compound having the formula (VI) with an amine having the formula R^4NH_2 to form a compound having the formula (VII):



where,

5 R^4 is a C_{3-15} cycloalkyl group, with or without one or more substituents, a C_{3-15} cycloalkenyl group, with or without one or more substituents, or a heterocycloalkyl group of 3 to 15 members, with or without one or more substituents; and

10 (v) removing the protecting portion of L, when L is the protected form of R^2 , on the compound having the formula (VII) to form the compound having the formula (I):



where,

R^2 is defined the same as R^1 above, with the proviso that at least one of R^1 and R^2 is not a hydrogen atom;

wherein, the one or more substituents are defined the same as for the one or more substituents of formula (I) above.

Pharmaceutically-Acceptable Dosage Forms

5 The compounds of the present invention may be administered to humans or other mammals by a variety of routes, including oral dosage forms and injections (intravenous, intramuscular, intraperitoneal, subcutaneous, and the like). Numerous other dosage forms containing the compounds of the present invention can be readily formulated by one skilled in the art, utilizing
10 the suitable pharmaceutical excipients as defined below. For considerations of patient compliance, oral dosage forms are generally most preferred.

 The rate of systemic delivery can be satisfactorily controlled by one skilled in the art, by manipulating any one or more of the following:

 (a) the active ingredient proper;

15 (b) the pharmaceutically-acceptable excipient(s), so long as the variants do not interfere in the activity of the particular active ingredient selected;

 (c) the type of excipient(s), and the concomitant desirable thickness and permeability (swelling properties) of the excipient(s);

20 (d) the time-dependent conditions of the excipient(s);

 (e) the particle size of the granulated active ingredient; and

 (f) the pH-dependent conditions of the excipient(s).

 Pharmaceutically-acceptable excipients include flavoring agents, pharmaceutical-grade dyes or pigments, solvents, co-solvents, buffer

systems, surfactants, preservatives, sweetener agents, viscosity agents, fillers, lubricants, glidants, disintegrants, binders and resins.

Conventional flavoring agents may be used, such as those described in *Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., pp. 1288-1300 (1990), which is incorporated in its entirety by reference herein. The pharmaceutical compositions of the invention generally contain from about 0 to 2 % of flavoring agents.

Conventional dyes and/or pigments may also be used, such as those described in the *Handbook of Pharmaceutical Excipients*, by the American Pharmaceutical Association & the Pharmaceutical Society of Great Britain, pp. 81-90 (1986), which is incorporated in its entirety by reference herein. The pharmaceutical compositions of the invention generally contain from about 0 to 2 % of dyes and/or pigments.

The pharmaceutical compositions of the invention generally contain from about 0.1 to 99.9 % of solvent(s). A preferred solvent is water. Preferred co-solvents include ethanol, glycerin, propylene glycol, polyethylene glycol, and the like. The pharmaceutical compositions of the invention may include from about 0 to 50 % of co-solvents.

Preferred buffer systems include acetic, boric, carbonic, phosphoric, succinic, malic, tartaric, citric, acetic, benzoic, lactic, glyceric, gluconic, glutaric and glutamic acids and their sodium, potassium and ammonium salts. Particularly preferred buffers are phosphoric, tartaric, citric and acetic acids and salts thereof. The pharmaceutical compositions of the invention generally contain from about 0 to 5 % of a buffer.

Preferred surfactants include polyoxyethylene sorbitan fatty acid esters, polyoxyethylene monoalkyl ethers, sucrose monoesters and lanolin esters and ethers, alkyl sulfate salts and sodium, potassium and ammonium salts of fatty acids. The pharmaceutical compositions of the invention
5 generally contain from about 0 to 2 % of surfactants.

Preferred preservatives include phenol, alkyl esters of parahydroxybenzoic acid, o-phenylphenol benzoic acid and salts thereof, boric acid and salts thereof, sorbic acid and salts thereof, chlorobutanol, benzyl alcohol, thimerosal, phenylmercuric acetate and nitrate, nitromersol,
10 benzalkonium chloride, cetylpyridinium chloride, methyl paraben and propyl paraben. Particularly preferred preservatives are the salts of benzoic acid, cetylpyridinium chloride, methyl paraben and propyl paraben. The pharmaceutical compositions of the invention generally include from about 0 to 2 % of preservatives.

15 Preferred sweeteners include sucrose, glucose, saccharin, sorbitol, mannitol and aspartame. Particularly preferred sweeteners are sucrose and saccharin. Pharmaceutical compositions of the invention generally include from about 0 to 5 % of sweeteners.

Preferred viscosity agents include methylcellulose, sodium
20 carboxymethylcellulose, hydroxypropyl-methylcellulose, hydroxypropylcellulose, sodium alginate, carbomer, povidone, acacia, guar gum, xanthan gum and tragacanth. Particularly preferred viscosity agents are methylcellulose, carbomer, xanthan gum, guar gum, povidone, sodium carboxymethylcellulose, and magnesium aluminum silicate. Pharmaceutical

compositions of the invention generally include from about 0 to 5 % of viscosity agents.

Preferred fillers include lactose, mannitol, sorbitol, tribasic calcium phosphate, diabasic calcium phosphate, compressible sugar, starch, calcium sulfate, dextro and microcrystalline cellulose. Pharmaceutical compositions of the invention generally contain from about 0 to 75 % of fillers.

Preferred lubricants/glidants include magnesium stearate, stearic acid and talc. Pharmaceutical compositions of the invention generally include from about 0 to 7 %, preferably, about 1 to 5 % of lubricants/glidants.

Preferred disintegrants include starch, sodium starch glycolate, crospovidone and croscarmellose sodium and microcrystalline cellulose. Pharmaceutical compositions of the invention generally include from about 0 to 20 %, preferably, about 4 to 15 % of disintegrants.

Preferred binders include acacia, tragacanth, hydroxypropylcellulose, pregelatinized starch, gelatin, povidone, hydroxypropylcellulose, hydroxypropylmethylcellulose, methylcellulose, sugar solutions, such as sucrose and sorbitol, and ethylcellulose. Pharmaceutical compositions of the invention generally include from about 0 to 12 %, preferably, about 1 to 10 % of binders.

Additional agents known to a skilled formulator may be combined with the compounds of the invention to create a single dosage form. Alternatively, additional agents may be separately administered to a mammal as part of a multiple dosage form.

For preparing pharmaceutical compositions containing the inventive compounds, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to 95 weight percent of active ingredient. Suitable solid carriers are known in the art, for example, magnesium carbonate, magnesium stearate, talc, sugar and lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically-acceptable carriers and methods of manufacture for various compositions may be found in *Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co. (1990), which is incorporated in its entirety by reference herein.

Liquid form preparations include solutions, suspensions and emulsions. Common liquid form preparations include water and water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation include solutions and solids in powder form, which may be combined with a pharmaceutically acceptable carrier, such as an inert compressed gas (e.g., nitrogen).

Also included are solid form preparations that may be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be delivered transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and emulsions and may be included in a transdermal patch of a matrix or reservoir type as is conventional in the art for this purpose.

5 The preferred mode of administering the compounds of the invention is oral. Preferably, the pharmaceutical preparation is in a unit dosage form. In such a form, the preparation is subdivided into suitable sized unit doses containing appropriate quantities of the active component, for example, an effective amount to achieve the desired purpose.

10 The quantity of active ingredient (compound) in a unit dose of preparation may be varied or adjusted from about 0.01 to 4,000 mg, preferably, from about 0.02 to 1,000 mg, more preferably, from about 0.3 to 500 mg, and most preferably, from about 0.04 to 250 mg, according to the particular application. A typical recommended daily dosage regimen for oral
15 administration can range from about 0.02 to 2,000 mg/day, in two to four divided doses. For convenience, the total daily dosage may be divided and administered in portions during the day as required. Typically, pharmaceutical compositions of the invention will be administered from about 1 to 5 times per day, or alternatively, as a continuous infusion. Such administration can be
20 used as a chronic or acute therapy. The amount of active ingredient that may be combined with carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5 to 95 % of active compound

(w/w). Preferably, such preparations will contain from about 20 to 80 wt. % of active compound.

The pharmaceutically-acceptable carriers employed in conjunction with the compounds of the present invention are used at a concentration sufficient to provide a practical size to dosage relationship. The pharmaceutically-acceptable carriers, in total, may comprise from about 0.1 to 99.9 % by weight of the pharmaceutical compositions of the invention, preferably, from about 20 to 80 % by weight.

Upon improvement of a patient's condition, a maintenance dose of a compound, composition or combination of the invention may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained. When the symptoms have been alleviated to the desired level, treatment should cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

Specific dosage and treatment regimens for any particular patient may be varied and will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health status, sex and diet of the patient, the time of administration, the rate of excretion, the specific drug combination, the severity and course of the symptoms being treated, the patient's disposition to the condition being treated and the judgment of the treating physician. Determination of the proper dosage regimen for a particular situation is within the skill of the art. The amount and

frequency of the administration of compounds of the invention or their pharmaceutically acceptable salts may be regulated according to the judgment of the attending clinician, based on the factors recited above. As a skilled artisan will appreciate, lower or higher doses than those recited above may be required.

For example, it is often the case that a proper dosage level is based on the weight of the patient. For instance, dosage levels of between about 0.01 and 100 mg/kg of body weight per day, preferably, between about 0.5 and 75 mg/kg of body weight per day, and more preferably, between about 1 and 50 mg/kg of body weight per day, of the inventive compounds, compositions and salts thereof described herein, are therapeutically useful for the treatment of a variety of biological disorders, particularly, male and female sexual dysfunction. Between two patients of differing weights, a higher dosage will be used for the heavier patient, all other things being equal.

The inventive compounds are understood to provide efficacious treatment of (male) erectile dysfunction, including a reasonable time of onset upon administration, and a reasonable duration after administration. For example, in the treatment of erectile dysfunction, a dosage of the inventive compound may be taken about an hour before a sex act is to be undertaken. Particular dosages will work within about thirty minutes of their administration. Ideal dosages will affect a patient within about fifteen minutes of their administration. While food, diet, pre-existing conditions, alcohol and other systemic conditions could lengthen the time delay for an inventive drug to work after its administration, it is understood that optimum dosages in

combination with sexual stimulation will result in an efficacious drug treatment within and for a reasonable amount of time.

The inventive compounds can exist in unsolvated as well as solvated forms, including hydrated forms. In general, the solvated forms, with
5 pharmaceutically- acceptable solvents, such as water, ethanol and the like, are equivalent to the unsolvated forms for purposes of this invention.

The inventive compounds may form pharmaceutically-acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic,
10 salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base forms with a sufficient amount of the desired acid to produce a salt in a conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base
15 solution, such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia or sodium bicarbonate. The free base forms may differ somewhat from their respective salt forms in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their respective free base forms for purposes of the invention.

20 The invention comprises a compound having the formula (I) or (II), a method for making an inventive compound, a method for making a pharmaceutical composition from at least one inventive compound and at least one pharmaceutically-acceptable carrier, and a method of using one or

more inventive compounds to treat a variety of disorders, symptoms and diseases.

The inventive compounds and their pharmaceutically-acceptable salt and neutral compositions may be formulated together with a pharmaceutically-acceptable carrier. The resulting composition may be administered *in vivo* to mammals, such as men or women, to treat a variety of disorders, symptoms and diseases. For example, the inventive compounds and compositions may be used to treat diseases of the urogenital system, specifically, male erectile dysfunction (e.g., impotence) and female sexual dysfunction. Male erectile dysfunction may be defined as an inability of the male to sufficiently obtain and/or sustain an erection to have intercourse with his mate. In the treatment of erectile dysfunction, it is believed that the inventive PDE V inhibitors of formulas (I) and (II) are beneficial therapeutic agents because they elevate cGMP levels in the human body. This action facilitates corpus cavernosum smooth muscle relaxation, which provides an increased flow of blood therein and results in an erection. This makes the inventive compounds especially useful for treating impotence and other types of diseases that are affected by cGMP levels.

Accordingly, another aspect of the invention is a method for treating erectile dysfunction in a mammal in need of such treatment, comprising administering to the mammal at least one compound having the formula (I) or (II) or a pharmaceutical composition thereof in an amount effective to ameliorate and/or reduce one or more of the symptoms associated with

erectile dysfunction sufficiently enough so that the mammal can complete intercourse with another mammal.

Introduced in 1998 as the first pill to treat impotence, Viagra[®] today is the most commonly prescribed medication to treat physiologically-caused
5 erectile dysfunction ("ED"). Certain patients, however, can experience undesirable side effects while taking Viagra[®]. For instance, the use of Viagra[®] is contraindicated to patients who are using organic nitrates, either regularly or intermittently. *Physicians' Desk Reference*[®], 55th Ed, pp. 2534-37 (2001). Combining Viagra[®] with nitrates can cause a hypotensive episode or suddenly
10 reduce blood pressure to dangerous levels, which may cause a heart attack. *Id.* Accordingly, men who have a heart condition that requires the use of nitrate drugs should not use Viagra[®]. *Id.* It has also been reported that Viagra[®] can cause a vision side effect by impairing the patient's color discrimination (blue/green), causing a "blue-halo" light visual alteration. *Id.*
15 This side effect is presumably due to inhibition of the PDE VI isoenzyme (found in a retina). *Id.*

An advantage of the inventive compounds is that they can be particularly selective for the PDE V isoenzyme in comparison to other types of PDE isoenzymes, such as the PDE VI isoenzyme. It is believed that this
20 increased selectivity will ameliorate side effects associated with the use of Viagra[®]. In particular, the high selectivity of the inventive compounds should minimize, and may even prevent, the occurrence of a "blue-halo" light visual alteration. It is believed that the increased isoenzyme selectivity in inhibiting

PDE V isoenzyme (found in a penis) versus PDE VI isoenzyme (found in a retina) accounts for obviating the "blue-halo" visual side effect.

Furthermore, a representative inventive compound did not adversely react with nitrate medication in a rat. It is believed the same lack of adverse interaction will apply to all inventive compounds in all mammals, including humans. An adverse reaction with nitrate medication may be dangerous and fatal. Adverse reactions include any reaction that could jeopardize or otherwise diminish the body's physiological functions. More specifically, in the case of combination therapy for a patient, comprising administering to the patient a nitrate donating agent combined with a PDE V inhibitor agent, an adverse nitrate reaction would be one in which the patient's blood pressure drops significantly more than with either agent administered alone.

This feature opens up a method of erectile dysfunction treatment to many patients who suffer from both an erectile dysfunction and a cardiovascular or other disease(s) that is treated with a nitrate donating medicament. Patients suffering from two or more different ailments that require dual (or multiple) treatments may have been born with one or both ailments, or later developed one or both ailments due to genetics or some other type of injury or disease, such as nerve damage, spinal cord injury, diabetes, and the like. Accordingly, it is another embodiment of this invention to treat a patient suffering from both (1) an erectile dysfunction and (2) at least one condition that can be treated with a nitrate donor medication, the inventive treatment comprising, a combination therapy comprising, an administration to a mammal of at least one inventive compound or a pharmaceutical

composition thereof, and at least one nitrate donating compound or a pharmaceutical composition thereof. The patient suffering from both erectile dysfunction and a need for a nitrate donating medicament can be treated for both conditions sequentially, concurrently and/or simultaneously. The combination therapy can be taken separately in any form, preferably in oral or patch doses, or can be formulated together for a single, combined dosage.

The compounds of the present invention may be employed alone or in combination with other agents, particularly, other types of PDE inhibitors (especially cGMP PDE V inhibitors), prostanoids, α -adrenergic receptor, dopamine receptor agonists, melanocortin receptor agonists, endothelin receptor antagonists, endothelin converting enzyme inhibitors, angiotensin II receptor antagonists, angiotensin converting enzyme inhibitors, neutral metalloendopeptidase inhibitors, renin inhibitors, serotonin 5-HT_{2c} receptor agonists, nociceptin receptor agonists, rho kinase inhibitors, potassium channel modulators and inhibitors of multidrug resistance protein 5.

Examples of therapeutic agents that may be used in combination with compounds of the invention are the following: PDE V inhibitors, such as sildenafil citrate (Viagra[®], Pfizer, Connecticut, United States), Vardenafil[™] (Bayer, Germany) and IC-351 (Cialis[™], Lilly-ICOS, Washington and Indiana, United States); prostanoids, such as prostaglandin E₁; α -adrenergic agonists, such as phentolamine mesylate; dopamine receptor agonists, such as apomorphine; angiotensin II antagonists, such as losartan, irbesartan, valsartan and candesartan; and ET_A antagonists, such as bosentan and ABT-627.

It is understood that other combinations may be undertaken while remaining within the scope of the invention. While one or more of the inventive compounds may be used in an application of monotherapy to treat erectile dysfunction, they also may be used in combination therapy, in which the inventive compounds are combined with one or more other pharmaceutical compounds that are useful for treating erectile dysfunction and/or other types of disorders, symptoms and diseases.

As discussed above, due to their cGMP-PDE V inhibitory activities, the inventive compounds are useful for treating urological disorders, in particular, female and male sexual dysfunctions. Other physiological disorders, symptoms and diseases can also benefit from cGMP-PDE V inhibition. More specifically, the inventive compounds, salts and derivatives thereof may be used to treat cardiovascular and cerebrovascular diseases, angina pectoris, hypertension, restenosis post angioplasty, endarterectomy, stent introduction, peripheral vascular diseases, cerebral stroke, respiratory tract disorders, such as reversible airway obstruction, chronic asthma and bronchitis, allergic disorders associated with atopy, such as urticaria, eczema, and rinitis, pulmonary hypertension, ischemic heart diseases, impaired glucose tolerance, diabetes and related complications, insulin resistance syndrome, hyperglycemia, polycystic ovarian syndrome, glomerular diseases, renal insufficiency, nephritis, tubular interstitial disease, autoimmune diseases, glaucoma, intestinal motility disorders, cachexia and cancer.

Another aspect of this invention is to provide a kit comprising separate containers in a single package, wherein the inventive pharmaceutical

compounds, compositions and/or salts thereof are used in combination with pharmaceutically-acceptable carriers to treat disorders, symptoms and diseases where cGMP-PDE V inhibition plays a role.

It is understood that other combinations may be undertaken while remaining within the scope of the invention. While one or more of the inventive compounds may be used in an application of monotherapy to treat erectile dysfunction, they also may be used in combination therapy, in which the inventive compounds are combined with one or more other pharmaceutical compounds that are useful for treating erectile dysfunction and/or other types of disorders, symptoms and diseases.

As discussed above, due to their cGMP-PDE V inhibitory activities, the inventive compounds are useful for treating urological disorders, in particular, female and male sexual dysfunctions. Other physiological disorders, symptoms and diseases can also benefit from cGMP-PDE V inhibition. More specifically, the inventive compounds, salts and derivatives thereof may be used to treat cardiovascular and cerebrovascular diseases, angina pectoris, hypertension, restenosis post angioplasty, endarterectomy, stent introduction, peripheral vascular diseases, cerebral stroke, respiratory tract disorders, such as reversible airway obstruction, chronic asthma and bronchitis, allergic disorders associated with atopy, such as urticaria, eczema, and rinitis, pulmonary hypertension, ischemic heart diseases, impaired glucose tolerance, diabetes and related complications, insulin resistance syndrome, hyperglycemia, polycystic ovarian syndrome, glomerular diseases, renal


insufficiency, nephritis, tubular interstitial disease, autoimmune diseases, glaucoma, intestinal motility disorders, cachexia and cancer.

Another aspect of this invention is to provide a kit comprising separate containers in a single package, wherein the inventive pharmaceutical compounds, compositions and/or salts thereof are used in combination with pharmaceutically-acceptable carriers to treat disorders, symptoms and diseases where cGMP-PDE V inhibition plays a role.

The above description is not intended to detail all modifications and variations of the invention. It will be appreciated by those skilled in the art that changes can be made to the embodiments described above without departing from the inventive concept. It is understood, therefore, that the invention is not limited to the particular embodiments described above, but is intended to cover modifications that are within the spirit and scope of the invention, as defined by the language of the following claims.

What is claimed is:

1. A compound, including an enantiomer, stereoisomer, rotomer, tautomer and/or prodrug thereof, or a pharmaceutical composition thereof, the compound having the formula (I):




(I)

where,

- (a) R^1 and R^2 are, independently of one another, each a C_{1-15} alkyl group, branched or straight chain, with or without one or more substituents, a C_{2-15} alkenyl group, branched or straight chain, with or without one or more substituents, a C_{2-15} alkynyl group, branched or straight chain, with or without one or more substituents, a C_{3-15} cycloalkyl group, with or without one or more substituents, an arylalkyl group, with or without one or more substituents, an aryl group, with or without one or more substituents, a heteroaryl group, with or without one or more substituents, $-OR^5$, $-COOR^5$, $-C(O)R^5$ or $-C(O)N(R^5)_2$, where, R^5 is a hydrogen atom or a hydrocarbon radical, with or without one or more substituents, or one of R^1 and R^2 is a hydrogen atom and the other one of R^1 and R^2 is defined the same as above;

- (b) R^3 is an aryl group, with or without one or more substituents, a heteroaryl group, with or without one or more substituents, or a heterocyclic group having 1 to 3 heteroatoms fused to a 5- or 6-membered aryl ring, with or without one or more substituents, with the proviso that R^3 is not an aryl group substituted at its para position with a -Y-aryl group, where, Y is a carbon-carbon single bond, -CO-, -O-, -S-, -N(R^{21})-, -CON(R^{22})-, -N(R^{22})CO-, -OCH₂-, -CH₂O-, -SCH₂-, -CH₂S-, -NHC(R^{23})(R^{24})-, -NR²³SO₂-, -SO₂NR²³-, -(R^{23})(R^{24})NH-, -CH=CH-, -CF=CF-, -CH=CF-, -CF=CH-, -CH₂CH₂-, -CF₂CF₂-,



where,

R^{21} is a hydrogen atom or a -CO(C₁₋₄ alkyl), C₁₋₆ alkyl, allyl, C₃₋₆ cycloalkyl, phenyl or benzyl group;

R^{22} is a hydrogen atom or a C₁₋₆ alkyl group;

R^{23} is a hydrogen atom or a C₁₋₅ alkyl, aryl or -CH₂-aryl group;


R^{24} is a hydrogen atom or a C₁₋₄ alkyl group;

R^{25} is a hydrogen atom or a C₁₋₈ alkyl, C₁₋₈ perfluoroalkyl,

C₃₋₆ cycloalkyl, phenyl or benzyl group;

R²⁶ is a hydrogen atom or a C₁₋₆ alkyl, C₃₋₆ cycloalkyl, phenyl or benzyl group;

R²⁷ is -NR²³R²⁴, -OR²⁴, -NHCONH₂, -NHCSNH₂,



and


R²⁸ and R²⁹ are, independently of one another, each a C₁₋₄ alkyl group or, taken together with each other, a -(CH₂)_q group, where q is 2 or 3; and

10 (c) R⁴ is a C₃₋₁₅ cycloalkyl group, with or without one or more substituents, a C₃₋₁₅ cycloalkenyl group, with or without one or more substituents, or a heterocycloalkyl group of 3 to 15 members, with or without one or more substituents;

15 wherein, the one or more substituents for all the groups are chemically-compatible and are, independently of one another, each an: alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, arylalkyl, alkylaryl, aryl, heteroaryl, heterocycloalkyl, hydroxyalkyl, arylalkyl, aminoalkyl, haloalkyl, thioalkyl, alkylthioalkyl, carboxyalkyl, imidazolylalkyl, indolylalkyl, mono-, di- and trihaloalkyl, mono-, di- and trihaloalkoxy, amino, alkylamino, dialkylamino, 20 alkoxy, hydroxy, halo, nitro, oximino, -COOR⁵⁰, -COR⁵⁰, -SO_{0.2}R⁵⁰, -SO₂NR⁵⁰R⁵¹, NR⁵²SO₂R⁵⁰, =C(R⁵⁰R⁵¹), =N-OR⁵⁰, =N-CN, =C(halo)₂, =S, =O,

-CON(R⁵⁰R⁵¹), -OCOR⁵⁰, -OCON(R⁵⁰R⁵¹), -N(R⁵²)CO(R⁵⁰), -N(R⁵²)COOR⁵⁰ or -N(R⁵²)CON(R⁵⁰R⁵¹) group, where:

5 R⁵⁰, R⁵¹ and R⁵² are, independently of one another, each a hydrogen atom or a branched or straight-chain, optionally substituted, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₄₋₆ heterocycloalkyl, heteroaryl or aryl group, or R⁵⁰ and R⁵¹ are joined together to form a carbocyclic or heterocyclic ring system, or R⁵⁰, R⁵¹ and R⁵² are, independently of one another, each:



where,

R^{40} and R^{41} are, independently of one another, each a hydrogen atom or a branched or straight-chain, optionally substituted, alkyl, cycloalkyl, heterocycloalkyl, halo, aryl, imidazolylalkyl, indolylalkyl,

heteroaryl, arylalkyl, arylalkoxy, heteroarylalkyl, heteroarylalkoxy, aminoalkyl, haloalkyl, mono-, di- or trihaloalkyl, mono-, di- or trihaloalkoxy, nitro, cyano, alkoxy, hydroxy, amino, phosphino, phosphate, alkylamino, dialkylamino, formyl, alkylthio, trialkylsilyl, alkylsulfonyl, arylsulfonyl, alkylsulfinyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, hydroxyalkyl, morpholino, thioalkyl, alkylthioalkyl, carboxyalkyl, oximino, $-\text{COOR}^{50}$, $-\text{COR}^{50}$, $-\text{SO}_{0-2}\text{R}^{50}$, $-\text{SO}_2\text{NR}^{50}\text{R}^{51}$, $-\text{NR}^{52}\text{SO}_2\text{R}^{50}$, $-\text{CON}(\text{R}^{50}\text{R}^{51})$, $-\text{OCON}(\text{R}^{50}\text{R}^{51})$, $-\text{N}(\text{R}^{52})\text{CO}(\text{R}^{50})$, $-\text{N}(\text{R}^{52})\text{COOR}^{50}$, $-\text{N}(\text{R}^{52})\text{CON}(\text{R}^{50}\text{R}^{51})$ or $-\text{OCONR}^{50}$ group, where, R^{50} , R^{51} and R^{52} are defined the same as above;

R^{42} is a hydrogen atom or a branched or straight-chain, optionally substituted, alkyl, alkenyl, arylalkyl or acyl group; and

R^{43} is a hydrogen atom or a branched or straight-chain, optionally substituted, alkyl or aryl group;

wherein, the optional substituents are defined the same as above for the one or more substituents.

2. The compound or pharmaceutical composition according to claim 1, where, R^1 is an alkyl or aryl group, with or without the one or more substituents.

3. The compound or pharmaceutical composition according to claim 2, where, R^1 is a methyl, ethyl or benzyl group, with or without the one or more substituents.

4. The compound or pharmaceutical composition according to claim 1, where, R^2 is an alkyl group, with or without the one or more substituents.

5

5. The compound or pharmaceutical composition according to claim 4, where, R^2 is a methyl, ethyl, iso-butyl or hydroxyethyl group, with or without the one or more substituents.

10

6. The compound or pharmaceutical composition according to claim 1, where, R^3 is an aryl group, with or without the one or more substituents.

15

7. The compound or pharmaceutical composition according to claim 6, where, R^3 is a hydroxyaryl, alkoxyaryl or aminosulfonylaryl group, with or without the one or more substituents.

20

8. The compound or pharmaceutical composition according to claim 7, where, the hydroxyaryl, alkoxyaryl or aminosulfonylaryl group for R^3 is substituted with at least one halogen atom on the aryl ring.
9. The compound or pharmaceutical composition according to claim 1, where, R^4 is a cycloalkyl or heterocycloalkyl group, with or without the one or more substituents.

10. The compound or pharmaceutical composition according to claim 9, where, R^4 is a cyclohexyl, hydroxycyclopentyl or tetrahydropyranyl group, with or without the one or more substituents.

5

11. The compound or pharmaceutical composition according to claim 1, where, R^1 is a methyl or ethyl group, R^2 is a methyl, ethyl or hydroxyethyl group, R^3 is a 3-chloro-4-hydroxyphenyl, 3-bromo-4-hydroxyphenyl, 3-chloro-4-methoxyphenyl, 3-bromo-4-methoxyphenyl, or 4-aminosulfonylphenyl group and R^4 is a cyclohexyl, tetrahydropyranyl or 2(R)-hydroxy-1(R)-cyclopentyl group.



10

12. The compound or pharmaceutical composition according to claim 1, where, R^1 is an alkyl or aryl group, with or without the one or more substituents, R^2 is an alkyl group, with or without the one or more substituents, and R^3 is a 4-hydroxyphenyl, 3-chloro-4-hydroxyphenyl, 3-bromo-4-hydroxyphenyl, 4-methoxyphenyl, 3-chloro-4-methoxyphenyl, 3-bromo-4-methoxyphenyl, 4-aminosulfonylphenyl, 3-chloro-4-aminosulfonylphenyl or 3-bromo-4-aminosulfonylphenyl group.



15

13. The compound or pharmaceutical composition according to claim 1,
which is:

5





10





15


20




5



10




or




15

14. The compound or pharmaceutical composition according to claim 1,
which is:


20




15. The compound or pharmaceutical composition according to claim 1,
which is:




16. The compound or pharmaceutical composition according to claim 1,
which is:




17. The compound or pharmaceutical composition according to claim 1,
which is:




18. The compound or pharmaceutical composition according to claim 1,
which is:




- 10 19. The compound or pharmaceutical composition according to claim 1,
which is:




20. The compound or pharmaceutical composition according to claim 1,
which is:




21. The compound or pharmaceutical composition according to claim 1,
which is:



22. The compound or pharmaceutical composition according to claim 1,
which is:



23. The compound or pharmaceutical composition according to claim 1,
which is:




24. The compound or pharmaceutical composition according to claim 1,
which has a PDE V IC₅₀ within the range of up to about 5 nM.

5 25. The compound or pharmaceutical composition according to claim 1,
which has a ratio of PDE VI IC₅₀/ PDE V IC₅₀ of > about 140.

26. The compound or pharmaceutical composition according to claim 1,
which has a PDE V IC₅₀ within the range of up to about 5 nM and a
10 ratio of PDE VI IC₅₀/ PDE V IC₅₀ of > about 140.

27. The compound or pharmaceutical composition according to claim 1,
where, R⁴ is:



15 where,

R¹, R² and R³, independently of one another, are each defined
the same as above for the compound of formula (I);

R⁹ is a hydrogen atom or an optionally substituted, oximino,
carboxyalkyl, C₁₋₆ alkoxy C₁₋₆ alkyl group, aryloxy C₁₋₆ alkyl, C₃₋₆
20 cycloalkoxy C₁₋₆ alkyl, heteroaryloxy C₁₋₆ alkyl, -COOH, ester, C₁₋₆

alkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocyclic, hydroxy C₁₋₆ alkyl, aryl or heteroaryl group;

R¹⁰ and R¹¹ are substituents on the same or different carbon atoms of the ring and, independently of one another, are each:

- 5 (a) defined the same as above for R⁹;
- (b) a hydroxy group or an ester group derived from a hydroxy group with a (i) C₁₋₆ carboxylic acid; (ii) C₃₋₆ cycloalkyl C₁₋₆ carboxylic acid; (iii) aryl C₁₋₆ carboxylic acid; or (iv) heteroaryl C₁₋₆ carboxylic acid group; or
- 10 (c) a C₁₋₆ alkoxy, amino, C₁₋₆ mono- or dialkylamino, C₁₋₆ alkylacylamino, C₁₋₆ alkylsulfonylamino or -NHCON(R¹⁴)₂ group, with or without one or more substituents, where R¹⁴ is a hydrogen atom or an
- 15 optionally substituted, alkyl or aryl group, or

R¹⁰ and R¹¹, taken together with each other and, optionally, with one or more carbon and/or hetero atoms of the ring, form an optionally substituted, spiro- or linearly fused, bi- or tri-cyclic ring system of from 8 to 12 members, including from 0 to 4 hetero atoms;

20 m and n, independently of one other, are each from 1 to 3; and

X is a chemically-compatible group, which is -C(R¹⁰R¹¹)-, -S(O)_y, -O- -N(R⁶⁰)-, where:

R¹⁰ and R¹¹ are defined the same as above;

y is from 0 to 2; and

R^{60} is a hydrogen atom or an optionally substituted, C_{1-8} alkyl, C_{1-8} alkynyl, C_{1-8} alkenyl, C_{3-8} cycloalkyl, aryl, heteroaryl, C_{4-8} heterocycloalkyl, COR^{61} , SO_2R^{61} , $COOR^{61}$, $CONR^{61}R^{62}$ or $SO_2NR^{61}R^{62}$ group, where:

5 R^{61} is a hydrogen atom or an optionally substituted, C_{1-8} alkyl, C_{2-8} alkynyl, C_{2-8} alkenyl, C_{3-8} cycloalkyl, aryl, heteroaryl or C_{4-8} heterocyclic group; and

R^{62} is a hydrogen atom or an optionally substituted, C_{1-8} alkyl, C_{2-8} alkynyl, C_{2-8} alkenyl, C_{3-8}

10 cycloalkyl, aryl, heteroaryl or C_{4-8} heterocyclic group; and

when R^{61} and R^{62} are the same or different alkyl groups, they are, optionally, joined together to form a carbocyclic or heterocyclic ring system;

15 wherein, the optional substituents are defined the same as for the one or more substituents of formula (I) above.

28. The compound or pharmaceutical composition according to claim 27, where, R^3 is an optionally substituted, hydroxyaryl, alkoxyaryl or aminosulfonylaryl group, wherein, the optional substituents are defined

20 the same as for the one or more substituents of formula (I) above.

29. The compound or pharmaceutical composition according to claim 27, where, R^9 is a hydrogen atom.

30. The compound or pharmaceutical composition according to claim 27, where, one of R^{10} and R^{11} is a hydrogen atom, and the other one of R^{10} and R^{11} is a hydrogen atom or a hydroxy group.

5 31. A method for treating a physiological disorder, symptom or disease in a patient, comprising administering to the patient an effective amount of the compound or pharmaceutical composition according to claim 1, wherein the physiological disorder, symptom or disease is urogenital, cardiovascular, cerebrovascular, peripheral vascular, angina pectoris, hypertension, restenosis post angioplasty, endarterectomy, stent
10 introduction, cerebral stroke, respiratory tract, allergic associated with atopy, pulmonary hypertension, ischemic heart, impaired glucose tolerance, diabetes and its related complications, insulin resistance syndrome, hyperglycemia, polycystic ovarian syndrome, glomerular, renal insufficiency, nephritis, tubular interstitial, autoimmune,
15 glaucoma, intestinal motility, cachexia or cancer.

32. The method according to claim 31, wherein the physiological disorder is a urogenital disorder.

20 33. The method according to claim 32, wherein the urogenital disorder is an erectile dysfunction.

34. A method for elevating a cGMP level in a patient in need of the treatment, comprising administering to the patient an effective amount of the compound or pharmaceutical composition according to claim 1.

5 35. A method for treating an erectile dysfunction in a patient in need of the treatment, comprising administering to the patient an effective amount of at least one of the compound or pharmaceutical composition according to claim 1.

10 36. The method of claim 35, wherein the patient has been, is being and/or will be treated with a nitrate donating pharmaceutical composition.


15 37. A method for treating an erectile dysfunction in a patient in need of the treatment, comprising administering to the patient an effective amount of at least one of the compound or pharmaceutical composition according to claim 27.

38. The method of claim 37, wherein the patient has been, is being and/or will be treated with a nitrate donating pharmaceutical composition.

20 39. A method for treating an erectile dysfunction and/or another symptom, disease or disorder in a patient in need of the treatment, comprising administering to the patient a combination therapy, comprising an effective amount of at least one of the compound or pharmaceutical

composition according to claim 1 and at least one compound selected from the group consisting of: a prostanoid, α -adrenergic receptor, dopamine receptor agonist, melanocortin receptor agonist, endothelin receptor antagonist, endothelin converting enzyme inhibitor, angiotensin II receptor antagonist, angiotensin converting enzyme inhibitor, neutral metalloendopeptidase inhibitor, renin inhibitor, serotonin 5-HT_{2c} receptor agonist, nociceptin receptor agonist, rho kinase inhibitor, potassium channel modulator and multidrug resistance protein 5 inhibitor.

40. A method for producing a compound having the formula (I), comprising:
- (i) reacting a compound having the formula (III) with an alkyl halide in the presence of a base to form a compound having the formula (IV):



where,


- (a) R¹ is a hydrogen atom or a C₁₋₁₅ alkyl group, branched or straight chain, with or without one or more substituents, a C₂₋₁₅ alkenyl group, branched or straight chain, with or without one or more substituents, a C₂₋₁₅ alkynyl group, branched or straight chain,

with or without one or more substituents, a C₃₋₁₅ cycloalkyl group,
 with or without one or more substituents, an arylalkyl group, with or
 without one or more substituents, an aryl group, with or without one
 or more substituents, a heteroaryl group, with or without one or
 more substituents, -OR⁵, -COOR⁵, -C(O)R⁵ or -C(O)N(R⁵)₂, where
 R⁵ is a hydrogen atom or a hydrocarbon radical, branched or
 straight-chain, with or without one or more substituents;

(b) L is R² or a protected form of R²; and

(c) Ph is a phenyl group;

(ii) debenzylating and then alkylating the compound having
 the formula (IV) with an alkyl halide having the formula
 XCH₂R³ to form the compound having the formula (V):




where,

X is a halogen atom and

R³ is an aryl group, with or without one or more
 substituents, a heteroaryl group, with or without one or more

substituents, or a heterocyclic group having 1 to 3 heteroatoms fused to a 5- or 6-membered aryl ring, with or without one or more substituents, with the proviso that R^3 is not an aryl group substituted at its para position with a -Y-aryl group, where Y is a carbon-carbon single bond, -CO-, -O-, -S-, -N(R^{21})-, -CON(R^{22})-, -N(R^{22})CO-, -OCH₂-, -CH₂O-, -SCH₂-, -CH₂S-, -NHC(R^{23})(R^{24})-, -NR²³SO₂-, -SO₂NR²³-, -C(R^{23})(R^{24})NH-, -CH=CH-, -CF=CF-, -CH=CF-, -CF=CH-, -CH₂CH₂-, -CF₂CF₂-,



where,

R^{21} is a hydrogen atom or a -CO(C₁₋₄ alkyl), C₁₋₆ alkyl, allyl, C₃₋₆ cycloalkyl, phenyl or benzyl group;

R^{22} is a hydrogen atom or a C₁₋₆ alkyl group;

R^{23} is a hydrogen atom or a C₁₋₅ alkyl, aryl or -CH₂-aryl group;


R^{24} is a hydrogen atom or a C₁₋₄ alkyl group;

R^{25} is a hydrogen atom or a C₁₋₈ alkyl, C₁₋₈ perfluoroalkyl;

C₃₋₆ cycloalkyl, phenyl or benzyl group;

R^{26} is a hydrogen atom or a C_{1-6} alkyl, C_{3-6} cycloalkyl, phenyl or benzyl group;

R^{27} is $-NR^{23}R^{24}$, $-OR^{24}$, $-NHCONH_2$, $-NHCSNH_2$,



5

and

R^{28} and R^{29} are, independently of one another, each a C_{1-4} alkyl group, or R^{28} and R^{29} , taken together with each other, are a $-(CH_2)_q$ group, where q is 2 or 3;

wherein, R^{21} through R^{29} are optionally substituted with one or more substituents; and

10

- (iii) deprotonating and then halogenating the compound having the formula (V) to form a compound having the formula (VI):


15



where,

Hal is a halogen atom;


- (iv) reacting the compound having the formula (VI) with an amine having the formula R^4NH_2 to form a compound having the formula (VII):



where,

R^4 is a C_{3-15} cycloalkyl group, with or without one or more substituents, a C_{3-15} cycloalkenyl group, with or without one or more substituents, or a heterocycloalkyl group of 3 to 15 members, with or without one or more substituents; and

- (v) removing the protecting portion of L, when L is the protected form of R^2 , on the compound having the formula (VII) to form the compound having the formula (I):



where,

R^2 is defined the same as R^1 above, with the proviso that at least one of R^1 and R^2 is not a hydrogen atom;

5

wherein, the one or more substituents for all the groups are chemically-compatible and are, independently of one another, each an: alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, arylalkyl, alkylaryl, aryl, heteroaryl, heterocycloalkyl, hydroxyalkyl, arylalkyl, aminoalkyl, haloalkyl, thioalkyl, alkylthioalkyl, carboxyalkyl, imidazolylalkyl, indolylalkyl, mono-, di- and trihaloalkyl, mono-, di- and trihaloalkoxy, amino, alkylamino, dialkylamino, alkoxy, hydroxy, halo, nitro, oximino, $-\text{COOR}^{50}$, $-\text{COR}^{50}$, $-\text{SO}_{0.2}\text{R}^{50}$, $-\text{SO}_2\text{NR}^{50}\text{R}^{51}$, $\text{NR}^{52}\text{SO}_2\text{R}^{50}$, $=\text{C}(\text{R}^{50}\text{R}^{51})$, $=\text{N}-\text{OR}^{50}$, $=\text{N}-\text{CN}$, $=\text{C}(\text{halo})_2$, $=\text{S}$, $=\text{O}$, $-\text{CON}(\text{R}^{50}\text{R}^{51})$, $-\text{OCOR}^{50}$, $-\text{OCON}(\text{R}^{50}\text{R}^{51})$, $-\text{N}(\text{R}^{52})\text{CO}(\text{R}^{50})$, $-\text{N}(\text{R}^{52})\text{COOR}^{50}$ or $-\text{N}(\text{R}^{52})\text{CON}(\text{R}^{50}\text{R}^{51})$ group, where:


10

15

R^{50} , R^{51} and R^{52} are, independently of one another, each a hydrogen atom or a C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{4-6} heterocycloalkyl, heteroaryl and aryl group, or R^{50} and R^{51} are joined together to form a carbocyclic or heterocyclic ring system, or R^{50} , R^{51} and R^{52} are, independently of one another, each:

20

120



where,

5

R^{40} and R^{41} are, independently of one another, each a hydrogen atom or an alkyl, cycloalkyl, heterocycloalkyl, halo, aryl, imidazolylalkyl, indolylalkyl, heteroaryl, arylalkyl, arylalkoxy, heteroarylalkyl,

heteroarylalkoxy, aminoalkyl, haloalkyl, mono-, di- or trihaloalkyl, mono-, di- or trihaloalkoxy, nitro, cyano, alkoxy, hydroxy, amino, phosphino, phosphate, alkylamino, dialkylamino, formyl, alkylthio, trialkylsilyl, alkylsulfonyl, arylsulfonyl, alkylsulfinyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, hydroxyalkyl, morpholino, thioalkyl, alkylthioalkyl, carboxyalkyl, oximino, $-\text{COOR}^{50}$, $-\text{COR}^{50}$, $-\text{SO}_{0-2}\text{R}^{50}$, $-\text{SO}_2\text{NR}^{50}\text{R}^{51}$, $-\text{NR}^{52}\text{SO}_2\text{R}^{50}$, $-\text{CON}(\text{R}^{50}\text{R}^{51})$, $-\text{OCON}(\text{R}^{50}\text{R}^{51})$, $-\text{N}(\text{R}^{52})\text{CO}(\text{R}^{50})$, $-\text{N}(\text{R}^{52})\text{COOR}^{50}$, $-\text{N}(\text{R}^{52})\text{CON}(\text{R}^{50}\text{R}^{51})$ or $-\text{OCONR}^{50}$ group, where, R^{50} , R^{51} and R^{52} are defined the same as above;

R^{42} is a hydrogen atom or an alkyl, alkenyl, arylalkyl or acyl group; and

R^{43} is a hydrogen atom or an alkyl or aryl group;

where, R^{40} through R^{43} and R^{50} through R^{52} are, independently of one another, each optionally substituted with any one of the groups defined above for the one or more substituents.

INTERNATIONAL SEARCH REPORT

Inter nal Application No

PCT/US 01/28983

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D473/04 C07D473/06 C07D473/08 A61K31/522 A61P15/00
A61P9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 05176 A (BEECHAM GROUP PLC) 2 April 1992 (1992-04-02) cited in the application claims ---	1-40
A	WO 92 05175 A (BEECHAM GROUP PLC) 2 April 1992 (1992-04-02) cited in the application claims ---	1-40
A	US 5 409 934 A (SMITH DAVID G ET AL) 25 April 1995 (1995-04-25) cited in the application claims --- -/--	1-40



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

28 January 2002

Date of mailing of the international search report

15/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

Interl 1al Application No

PC1/US 01/28983

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 389 282 A (BEECHAM WUELFING GMBH & CO KG ;BEECHAM GROUP PLC (GB)) 26 September 1990 (1990-09-26) claims ----	1-40
A	WO 99 62905 A (ALMIRALL PRODESFARMA SA ;GRACIA FERRER JORDI (ES); PRIETO SOTO JOS) 9 December 1999 (1999-12-09) claims ----	1,31,40
A	WO 94 28902 A (PFIZER LTD ;PFIZER (US); PFIZER RES & DEV (IE); ELLIS PETER (GB);) 22 December 1994 (1994-12-22) cited in the application claims ----	1,31,40
A	WO 99 24433 A (NIEWOEHNER ULRICH ;HANING HELMUT (DE); SERNO PETER (DE); BAYER AG) 20 May 1999 (1999-05-20) cited in the application claims ----	1,31,40
A	T. KATSUSHIMA ET AL: "Structure-activity relationships of 8-cycloalkyl-1,3-dipropylxanthines as antagonists of adenosine receptors" JOURNAL OF MEDICINAL CHEMISTRY., vol. 33, no. 7, 1990, pages 1906-1910, XP000611869 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document ----	1,31
A	CHEMICAL ABSTRACTS, vol. 106, no. 13, 30 March 1987 (1987-03-30) Columbus, Ohio, US; abstract no. 95577, N.I. ROMANENKO ET AL: "Synthesis and biological activity of 3-methyl,7- or 8-alkyl-,7,8-dialkyl, heterocyclic, and cyclohexylaminoxanthines" XP002188426 abstract & FARM. ZH., no. 5, 1986, pages 41-44, KIEV ----	1,31,40
A	FR 2 116 302 A (LABORATOIRE LEBRUN S.A.) 13 July 1972 (1972-07-13) claims -----	1,31,40

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/28983

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9205176	A	02-04-1992	AU 650679 B2	30-06-1994
			AU 8645891 A	15-04-1992
			CA 2092429 A1	27-03-1992
			EP 0550559 A1	14-07-1993
			WO 9205176 A1	02-04-1992
			IE 913351 A1	08-04-1992
			JP 6501691 T	24-02-1994
			MX 9101235 A1	04-05-1992
			NZ 239920 A	27-06-1994
			PT 99037 A	31-08-1992
			ZA 9107611 A	30-12-1992
WO 9205175	A	02-04-1992	AU 653364 B2	29-09-1994
			AU 8541391 A	15-04-1992
			CA 2092430 A1	27-03-1992
			EP 0550570 A1	14-07-1993
			WO 9205175 A1	02-04-1992
			IE 913350 A1	08-04-1992
			JP 6501251 T	10-02-1994
			MX 9101237 A1	04-05-1992
			NZ 239921 A	23-12-1993
			PT 99062 A	31-08-1992
			ZA 9107610 A	30-09-1992
US 5409934	A	25-04-1995	AT 194345 T	15-07-2000
			AU 649091 B2	12-05-1994
			AU 9122791 A	22-07-1992
			CA 2098846 A1	22-06-1992
			DE 69132291 D1	10-08-2000
			DE 69132291 T2	14-12-2000
			EP 0563168 A1	06-10-1993
			ES 2148170 T3	16-10-2000
			WO 9211260 A1	09-07-1992
			IE 914502 A1	01-07-1992
			JP 6504534 T	26-05-1994
			MX 9102766 A1	01-09-1992
			NZ 241158 A	22-12-1994
			PT 99919 A , B	31-12-1992
			ZA 9110058 A	15-04-1993
EP 0389282	A	26-09-1990	AU 629315 B2	01-10-1992
			AU 5208390 A	27-09-1990
			CA 2012686 A1	23-09-1990
			CN 1047502 A , B	05-12-1990
			CZ 9103918 A3	17-11-1993
			EP 1120417 A2	01-08-2001
			EP 0389282 A2	26-09-1990
			FI 95033 B	31-08-1995
			HU 54155 A2	28-01-1991
			IL 93832 A	31-07-1994
			JP 2273676 A	08-11-1990
			JP 2510889 B2	26-06-1996
			KR 160768 B1	01-12-1998
			NO 175592 B	25-07-1994
			NZ 233021 A	23-12-1993
			PL 164811 B1	31-10-1994
			PT 93527 A , B	07-11-1990
			SK 391891 A3	11-12-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/28983

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0389282 A		US 6180791 B1 US 5981535 A US 5734051 A ZA 9002170 A ZW 3690 A1	30-01-2001 09-11-1999 31-03-1998 27-03-1991 06-02-1991
WO 9962905 A	09-12-1999	AU 4501199 A WO 9962905 A1	20-12-1999 09-12-1999
WO 9428902 A	22-12-1994	AT 163852 T AU 676571 B2 AU 6797394 A CA 2163446 A1 CN 1124926 A ,B CZ 9503242 A3 DE 69408981 D1 DE 69408981 T2 DK 702555 T3 WO 9428902 A1 EP 0702555 A1 ES 2113656 T3 FI 955911 A GR 3026520 T3 IL 109873 A IL 121836 A JP 11286444 A JP 11263728 A JP 2925034 B2 JP 9503996 T KR 262926 B1 LV 12269 A LV 12269 B NO 954757 A NO 20000702 A NO 20000703 A NZ 266463 A PL 311948 A1 RU 2130776 C1 ZA 9404018 A	15-03-1998 13-03-1997 03-01-1995 22-12-1994 19-06-1996 17-07-1996 16-04-1998 02-07-1998 06-04-1998 22-12-1994 27-03-1996 01-05-1998 08-12-1995 31-07-1998 27-12-1998 27-12-1998 19-10-1999 28-09-1999 26-07-1999 22-04-1997 01-09-2000 20-05-1999 20-08-1999 24-11-1995 24-11-1995 24-11-1995 24-03-1997 18-03-1996 27-05-1999 08-12-1995
WO 9924433 A	20-05-1999	DE 19750085 A1 DE 19812462 A1 DE 19840289 A1 AU 738675 B2 AU 1558799 A BG 104406 A BR 9812785 A CN 1278822 T DE 19881732 D2 DK 200000766 A EE 200000291 A WO 9924433 A1 EP 1174431 A2 EP 1049695 A1 FI 20001086 A GB 2346877 A ,B HR 20000292 A1 HU 0100394 A2	20-05-1999 30-09-1999 09-03-2000 20-09-2001 31-05-1999 31-08-2001 10-10-2000 03-01-2001 24-08-2000 09-05-2000 15-06-2001 20-05-1999 23-01-2002 08-11-2000 09-05-2000 23-08-2000 30-04-2001 28-09-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter: International Application No

PCT/US 01/28983

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9924433 A		JP 2001522851 T LU 90561 A1 NO 20002444 A PL 340400 A1 SE 0001745 A SK 7092000 A3 TR 200001338 T2 ZA 9810297 A	20-11-2001 01-12-2000 11-05-2000 29-01-2001 11-05-2000 12-03-2001 21-08-2000 20-05-1999
FR 2116302 A	13-07-1972	FR 2116302 A5	13-07-1972